Vol. 26

Index/Contents

2009

Indexes

DOI: 10.1111/j.1464-5491.2009.02883.x

Author Index

Abbasi, M. *see* Esteghamati, A.
Abbasi, M. *see* Esteghamati, A.
Abdous, B. *see* Ouhoummane, N.
Abid, M. *see* Kamoun, M.
Abreu, M. *see* Chaves-Fonseca, R.M.
Agarwal, M.M., Dhatt, G.S., Othman, Y. & Gupta, R. Gestational diabetes: fasting capillary glucose as a screening test in a multi-ethnic, high-risk population, 760
Ahlquist, J. *see* Cooke, D.
Ahluwalia, R., Perkins, K., Ewins, D. & Goenka, N. Exenatide—a potential role in treatment of HNF1-alpha MODY in obese patients?, 834
Ahmed, P. *see* Choudhury, S.M.
Ahn, C.W. *see* Kim, H.C.
Ahn, C.W. *see* Park, S.E.
Ahn, S.A. *see* Hermans, M.P.
Akiba, T. *see* Hayashino, Y.
Akimoto, K. *see* Inoue, K.
Akinci, B. *see* Bayraktar, F.
Akizawa, T. *see* Hayashino, Y.
Akram, K., Pedersen-Bjergaard, U., Carstensen, B., Borch-Johnsen, K. & Thorsteinsson, B. Prospective and retrospective recording of severe hypoglycaemia, and assessment of hypoglycaemia awareness in insulin-treated Type 2 diabetes, 1308
Al-Ansari, S.A., Tennant, M.T.S., Greve, M.D.J., Hinz, B.J. & Senior, P.A. Short report: suboptimal diabetes care in high-risk diabetic patients attending a specialist retina clinic, 1296
Al-Bermani, A. *see* Arun, C.S.
Alberti, K.G.M.M. *see* Cameron, A.J.
Alberti, G. *see* Gao, W.G.
Aldasouqi, S.A. & Gossain, V.V. A proposal for a role of HbA_{1c} in screening for gestational diabetes, 833
Algert, C.S., McElduff, A., Morris, J.M. & Roberts, C.L. Perinatal risk factors for early onset of Type 1 diabetes in a 2000–2005 birth cohort, 1193
Allan, B. *see* Ng, J.M.
Allen, D. & Gregory, J. The transition from children's to adult diabetes services: understanding the 'problem', 162
Allet, L., Armand, S., de Bie, R., Golay, A., Pataky, Z., Aminian, K. & de Bruin, E. Clinical factors associated with gait alterations in diabetic patients, 1003
Alonso, M. *see* Martín-Frías, M.
Alte, D. *see* Schipf, S.
Alvarsson, M., Hilding, A. & Östenson, C.-G. Factors determining normalization of glucose intolerance in middle-aged Swedish men and women: a 8–10-year follow-up, 345
Amann-Zalan, I. *see* Nielsen, J.K.
Amiel, S.A. *see* Choudhary, P.
Amiel, S.A. *see* Rogers, H.
Amin, R. on behalf of the North West England Paediatric Diabetes Network. *see* Sun, S.
Aminian, K. *see* Allet, L.
Anantharaman, R. *see* Jayaprakash, P.
Anderson, J.M., Savvidou, M.D., Kaihura, C., McEniery, C.M. & Nicolaides, K.H. Maternal arterial stiffness in pregnancies affected by Type 1 diabetes mellitus, 1135
Andersson, S.W. *see* Ning, F.
Anderwald, C. *see* Stadler, M.
Andrianakos, A. *see* Tentolouris, N.
Angus, E. *see* Garg, R.
Apelqvist, J. *see* Owings, T.M.
Aragón-Sánchez, J. & Lázaro-Martínez, J.L. Comments on the use of bemiparin in diabetic foot ulcers, 110
Aragón-Sánchez, J., Lázaro-Martínez, J.L., Quintana-Marrero, Y., Hernández-Herrero, M.J., García-Morales, E., Cabrera-Galván, J.J. & Beneit-Montesinos, J.V. Are diabetic foot ulcers complicated by MRSA osteomyelitis associated with worse prognosis? Outcomes of a surgical series, 552
Arean, P. *see* Fisher, L.
Arellano, J. *see* Pratoomsoot, C.
Argente, J. *see* Rubio-Cabezas, O.
Arimura, K. *see* Deguchi, T.
Armand, S. *see* Allet, L.
Arun, C.S., Al-Bermani, A., Stannard, K. & Taylor, R. Long-term impact of retinal screening on significant diabetes-related visual impairment in the working age population, 489
Arutchelvam, V., Heise, T., Dellweg, S., Elbroend, B., Minns, I. & Home, P.D. Plasma glucose and hypoglycaemia following exercise in people with Type 1 diabetes: a comparison of three basal insulins, 1027
Asano, Y. *see* Hayashino, Y.
Asgari, F. *see* Esteghamati, A.
Ashraf, H. *see* Esteghamati, A.
Aso, Y. *see* Yamamoto-Suganuma, R.
Attia, J. *see* Lowe, J.M.
Auinger, M. *see* Stadler, M.
Avignon, A. *see* Radermecker, R.P.
Ayhan, E. *see* Onat, A.

Bachmann, M.O. *see* Misra, A.
Badet, L. *see* Benhamou, P.Y.
Baek, H.S. *see* Jin, H.Y.
Baertschiger, R. *see* Benhamou, P.Y.
Baik, S.H. *see* Yoo, H.J.
Bailey, C. *see* Patra, S.
Bakker, S.J.L. *see* Lambers Heerspink, H.J.
Balani, J., Hyer, S.L., Rodin, D.A. & Shehata, H. Pregnancy outcomes in women with gestational diabetes treated with metformin or insulin: a case–control study, 798
Balena, R. *see* Kapitza, C.
Balko, S.U. *see* Johnson, J.A.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Bansal, D., Bhansali, A., Hota, D., Chakrabarti, A. & Dutta, P. Amitriptyline vs. pregabalin in painful diabetic neuropathy: a randomized double blind clinical trial, 1019
Barakat, O. *see* Misra, A.
Barbetti, F. *see* Nocerino, V.
Barnard, K.D. *see* Speight, J.
Barnard, M. *see* Cooke, D.
Barnett, J. *see* Kirk, A.
Barrio, R. *see* Martin-Frias, M.
Bartley, P. *see* Moses, R.G.
Baskar, V. *see* Manley, S.E.
Baskar, V. *see* Manley, S.E.
Basu, A. *see* Ndip, A.
Basuyaux, O. *see* Dumont, I.J.
Baviera, G. *see* D'Anna, R.
Bayle, F. *see* Benhamou, P.Y.
Bayle, F. *see* Lablanche, S.
Bayraktar, F., Akinci, B., Demirkan, F., Yener, S., Yesil, S., Kırmaz, C. & Comlekci, A. Serum sickness-like reactions associated with type III insulin allergy responding to plasmapheresis, 659
Bebakar, W.M.W. *see* Marre, M.
Becker, M. *see* Owings, T.M.
Beckert, S., Sundermann, K., Wolf, S., Königsrainer, A. & Coerper, S. Haemodialysis is associated with changes in cutaneous microcirculation in diabetes mellitus, 89
Beer, N. *see* Edghill, E.L.
Befani, C.D. *see* Pikilidou, M.I.
Begg, L. *see* Burns, J.
Beijers, H.J.B.H., Losekoot, M., Odink, R.J. & Bravenboer, B. Hepatocyte nuclear factor (HNF)1A and HNF4A substitution occurring simultaneously in a family with maturity-onset diabetes of the young, 1172
Béliard, S., Nogueira, J.P., Maraninchi, M., Lairon, D., Nicolay, A., Giral, P., Portugal, H., Vialettes, B. & Valéro, R. Parallel increase of plasma apoproteins C-II and C-III in Type 2 diabetic patients, 736
Bell, P.M. *see* Cuthbertson, J.
Belloni, E. *see* Perseghin, G.
Beloosesky, Y. *see* Weiss, A.
Benbassat, J. & Polak, B.C.P. Reliability of screening methods for diabetic retinopathy, 783
Beneit-Montesinos, J.V. *see* Aragón-Sánchez, J.
Benhamou, P.Y., Milliat-Guittard, L., Wojtusciszyn, A., Kessler, L., Toso, C., Baertschiger, R., Debaty, I., Badet, L., Penfornis, A., Thivolet, C., Renard, E., Bayle, F., Morel, P., Morelon, E., Colin, C. & Berney, T. on behalf of the GRAGIL Group, Quality of life after islet transplantation: data from the GRAGIL 1 and 2 trials, 617
Benhamou, P.Y. *see* Lablanche, S.
Bergeron, J. *see* Ferland, A.
Berghold, A. *see* Siebenhofer, A.
Berney, T. on behalf of the GRAGIL Group *see* Benhamou, P.Y.
Bertoni, A.G. *see* Lutsey, P.L.
Bertrand, F. *see* Ferland, A.
Best, J.H., Boye, K.S., Rubin, R.R., Cao, D., Kim, T.H. & Peyrot, M. Improved treatment satisfaction and weight-related quality of life with exenatide once weekly or twice daily, 722
Bethel, M.A., Deedwania, P., Levitt, N.S., Schmitz, O., Huntsman-Labed, A., Califf, R.M., Haffner, S.M. & Diem, P. for the NAVIGATOR Study Group, Metabolic syndrome and alanine aminotransferase: a global perspective from the NAVIGATOR screening population, 1204
Betteridge, D.J. *see* Scheen, A.
Betteridge, D.J. *see* Scheen, A.J.
Bhagey, J. *see* Vernon, S.A.
Bhansali, A. *see* Bansal, D.
Bhansali, A. *see* Jayaprakash, P.
Bhansali, S. *see* Jayaprakash, P.
Bhartia, A. *see* Nagpal, J.
Bhushan, R. *see* Wolffenbuttel, B.H.R.
Bialasiewicz, P., Pawlowski, M., Nowak, D., Loba, J. & Czupryniak, L. Decreasing concentration of interstitial glucose in REM sleep in subjects with normal glucose tolerance, 339
de Bie, R. *see* Allet, L.
Bieglmayer, C. *see* Stadler, M.
Bilo, H.J.G., Logtenberg, S.J.J., Joosten, H., Groenier, K.H., Ubink-Veltmaat, L.J. & Kleefstra, N. Modification of diet in renal disease and Cockcroft–Gault formulas do not predict mortality (ZODIAC-6), 478
Birkeland, K. *see* Scheen, A.
Birkeland, K. *see* Scheen, A.J.
Birman, P. *see* Kapitza, C.
Bischof, M. *see* Stadler, M.
Bleyer, A.J., Hire, D., Russell, G.B., Xu, J., Divers, J., Shihabi, Z., Bowden, D.W. & Freedman, B.I. Ethnic variation in the correlation between random serum glucose concentration and glycated haemoglobin, 128
Boaz, M., Lisy, L., Zandman-Goddard, G. & Wainstein, J. The effect of anti-inflammatory (aspirin and/or statin) therapy on body weight in Type 2 diabetic individuals: EAT, a retrospective study, 708
Boaz, M. *see* Weiss, A.
Boehm, B. *see* Kempf, K.
Bogaty, P. *see* Ferland, A.
Boissonnet, C. *see* Escobedo, J.
Bolin, K., Gip, C., Mörk, A.-C. & Lindgren, B. Diabetes, healthcare cost and loss of productivity in Sweden 1987 and 2005—a register-based approach, 928
Bonfanti, R. *see* Nocerino, V.
Bonomo, M. *see* Lepore, G.
Booth, G. *see* Gucciardi, E.
Boraik, M. *see* Vernon, S.A.
Borch-Johnsen, K. *see* Akram, K.
Borch-Johnsen, K. *see* Dalsgaard, E.-M.
Borch-Johnsen, K. *see* Færch, K.
Borretta, G. *see* Tassone, F.
Boucher, B.J. Renal failure and rhabdomyolysis associated with sitagliptin and simvastatin use. But what about the amiodarone?, 192
Boulton, A.J.M. *see* Bowling, F.L.
Boulton, A.J.M. *see* Ndip, A.
Bowden, D.W. *see* Bleyer, A.J.
Bowling, F.L., King, L., Fadavi, H., Paterson, J.A., Preece, K., Daniel, R.W., Matthews, D.J. & Boulton, A.J.M. An assessment of the accuracy and usability of a novel optical wound measurement system, 93

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Boyages, S.C. *see* Mitchell, P.
Boye, K.S. *see* Best, J.H.
Boye, K.S. *see* Pratoomsoot, C.
Boye, K.S. *see* Zhao, Y.
Brackenridge, A.L., Jackson, N., Jefferson, W., Stolinski, M., Shojaee-Moradie, F., Hovorka, R., Umpleby, A.M. & Russell-Jones, D. Effects of rosiglitazone and pioglitazone on lipoprotein metabolism in patients with Type 2 diabetes and normal lipids, 532
Bradley, C. *see* Brose, L.S.
Bradley, C. *see* Sampson, M.J.
Bradley, C. Psychometric tools for measuring outcomes of diabetes education; a critique of Eigenmann *et al.*'s assessment of suitability, 952
Bramness, J.G. *see* Skurtveit, S.
Brändle, M. *see* Marre, M.
Brassard, P. *see* Ferland, A.
Bravenboer, B. *see* Beijers, H.J.B.H.
Brezis, M. *see* Stern, Z.
Bril, V., Tomioka, S., Buchanan, R.A., Perkins, B.A. the mTCNS. Reliability and validity of the modified Toronto Clinical Neuropathy Score in diabetic sensorimotor polyneuropathy, 240
Brinkman, J.W. *see* Lambers Heerspink, H.J.
Bristow, I.R. & Spruce, M.C. Fungal foot infection, cellulitis and diabetes: a review, 548
Brooks, A.M.S. & Lissett, C.A. A dramatic deterioration in diabetic retinopathy with improvement in glycated haemoglobin (HbA_{1c}) on exenatide treatment, 190
Brophy, S. *see* Choudhury, S.M.
Brophy, S. *see* Choudhury, S.M.
Brose, L.S., Mitchell, J. & Bradley, C. Comments on Speight *et al.*'s 'Not all roads lead to Rome—a review of quality of life measurement in adults with diabetes', 1076
Broshtilova, V., Ramot, Y., Molho-Pessach, V. & Zlotogorski, A. Diabetes mellitus may be the earliest and sole manifestation of the H syndrome, 1179
Bruce, J. *see* Hill, N.R.
de Bruin, E. *see* Allet, L.
Bruttomesso, D. *see* Lapolla, A.
Bruttomesso, D. *see* Lepore, G.
Buchanan, R.A. *see* Bril, V.
Buitrón, L.V. *see* Escobedo, J.
Bulpitt, C.J. *see* Hewitt, J.
Bulugahapitiya, U., Siyambalapitiya, S., Sithole, J. & Idris, I. Is diabetes a coronary risk equivalent? Systematic review and meta-analysis, 142
Burns, J., Wegener, C., Begg, L., Vicaretti, M. & Fletcher, J. Randomized trial of custom orthoses and footwear on foot pain and plantar pressure in diabetic peripheral arterial disease, 893
Bus, S.A. *see* Owings, T.M.
Byrne, C.D. *see* Turzyniecka, M.

Cabrera-Galván, J.J. *see* Aragón-Sánchez, J.
Cacciotti, L. *see* Spallone, V.
Cade, J.E., Kirk, S.F.L., Nelson, P., Hollins, L., Deakin, T., Greenwood, D.C. & Harvey, E.L. Can peer educators influence healthy eating in people with diabetes? Results of a randomized controlled trial, 1048
Calderon-Margalit, R. *see* Stern, Z.
Califf, R.M. *see* Bethel, M.A.
Cameron, A.J., Zimmet, P.Z., Shaw, J.E. & Alberti, K.G.M.M. The metabolic syndrome: in need of a global mission statement, 306
Campbell, J. *see* Corbould, A.
Campbell, L.V. *see* Lee, P.
Can, G. *see* Onat, A.
Canu, T. *see* Perseghin, G.
Cao, D. *see* Best, J.H.
Carstensen, B. *see* Akram, K.
Casbard, A. *see* Cooke, D.
Cass, A.E.G. *see* Oliver, N.S.
Castano, L. *see* Garin, I.
Catanzariti, L., Faulks, K., Moon, L., Waters, A.-M., Flack, J. & Craig, M.E. Australia's National trends in the incidence of Type 1 diabetes in 0–14-year-olds, 2000–2006, 596
Caumo, A. *see* Perseghin, G.
Cavanagh, P.R. *see* Owings, T.M.
Cederholm, J. *see* Gudbjörnsdottir, S.
Celewicz, Z. *see* Ogonowski, J.
Cerdà, L. *see* Rullan, M.
Cha, B.S. *see* Park, S.E.
Chakrabarti, A. *see* Bansal, D.
Champagne, B.M. on behalf of the CARMELA Study Investigators. *see* Escobedo, J.
Chan, B.K.S. *see* Norris, S.L.
Chan, J., So, W., Ko, G., Tong, P., Yang, X., Ma, R., Kong, A., Wong, R., Le Coguiec, F., Tamesis, B., Wolthers, T., Lyubomirsky, G. & Chow, P. The Joint Asia Diabetes Evaluation (JADE) Program: a web-based program to translate evidence to clinical practice in Type 2 diabetes, 693
Chang, H.-W. *see* Chen, M.-C.
Chang, H.Y. *see* Li, C.L.
Charbonnel, B. on behalf of the PROactive investigators. *see* Scheen, A.
Charbonnel, B. on behalf of the PROactive Investigators, *see* Scheen, A.J.
Charfi, N. *see* Kamoun, M.
Chaudhry, A. *see* Greenfield, J.R.
Chaves-Fonseca, R.M., Matos, O.S., Lordelo, R.A., Abreu, M., Farias, M.G., Coutinho, J.F., Ribeiro, M.N., Matteoni-Athayde, L., Lessa, I., Pousada, J., Oliveira, M., Lopes, C., Strock, E. & Mazze, R. Implementation of a systematic approach to diabetes in primary care in Bahia, Brazil improves metabolic outcomes: PRODIBA—Programa de Interiorização da Assistência ao Diabetes na Bahia (Project for Dissemination of Diabetes Care in the State of Bahia), 286
Chen, C.-Y. *see* Chen, M.-C.
Chen, J.-F. *see* Chen, M.-C.
Chen, M.-C., Sheu, J.-J., Wang, P.-W., Chen, C.-Y., Kuo, M.-C., Hsieh, C.-J., Chen, J.-F. & Chang, H.-W. Complications impaired endothelial progenitor cell function in Type 2 diabetic patients with or without critical leg ischaemia: implication for impaired neovascularization in diabetes, 134
Chen, S.S.H., Jenkins, A.J. & Majewski, H. Elevated plasma prostaglandins and acetylated histone in monocytes in Type 1 diabetes patients, 182
Chen, W.-Q. *see* Zhang, C.-X.
Chen, X. *see* Yang, H.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Chen, Y.-M. *see* Zhang, C.-X.
Cheow, S.P. *see* Dalan, R.
Chester, P. *see* Patel, A.
Cheung, N.W. *see* Gan, R.M.
Chian, K.Y. *see* Dalan, R.
Chiarelli, F. *see* Marcovecchio, M.L.
Chipperfield, A.J. *see* Turzyniecka, M.
Chitson, P. *see* Gao, W.G.
Cho, G.J. *see* Yoo, H.J.
Cho, Y.J. *see* Kim, H.C.
Choi, D.S. *see* Yoo, H.J.
Choi, K.M. *see* Yoo, H.J.
Choudhary, P., Lonnen, K., Emery, C.J., MacDonald, I.A., MacLeod, K.M., Amiel, S.A. & Heller, S.R. Comparing hormonal and symptomatic responses to experimental hypoglycaemia in insulin- and sulphonylurea-treated Type 2 diabetes, 665
Choudhury, S.M., Brophy, S. & Williams, R. Understanding and beliefs of diabetes in the UK Bangladeshi population, 636
Choudhury, S.M., Brophy, S., Fareedi, M.A., Zaman, B., Ahmed, P. & Williams, R. Examining the effectiveness of a peer-led education programme for Type 2 diabetes and cardiovascular disease in a Bangladeshi population, 40
Choudhury, S.R. *see* Peter, R.
Chow, P. *see* Chan, J.
Christiansen, J.S. *see* Nielsen, J.K.
Christiansen, T. *see* Dalsgaard, E.-M.
Christoyannis, F. *see* Tentolouris, N.
Chubb, S.A.P. *see* Hamilton, E.J.
Chubb, S.A.P. *see* Schimke, K.
Chudleigh, R. *see* Peter, R.
Cieza, A. *see* Kirchberger, I.
Clough, G.F. *see* Turzyniecka, M.
Coenen, M. *see* Kirchberger, I.
Coerper, S. *see* Beckert, S.
Coffey, L., Gallagher, P., Horgan, O., Desmond, D. & MacLachlan, M. Psychosocial adjustment to diabetes-related lower limb amputation, 1063
Colagiuri, S. on behalf of the LEAD-1 SU Study Group *see* Marre, M.
Colagiuri, R. & Eigenmann, C.A. A national consensus on outcomes and indicators for diabetes patient education, 442
Colagiuri, R. *see* Eigenmann, C.A.
Colin, C. *see* Benhamou, P.Y.
Colino, E. *see* Martín-Frías, M.
Collier, A., Ghosh, S., Davidson, D.F. & Kilpatrick, E.S. HbA1c: 'The old order changeth', 573
Collins, C.E. *see* Smart, C.E.
Collins, M.M., Corcoran, P. & Perry, I.J. Anxiety and depression symptoms in patients with diabetes, 153
Colombo, C. *see* Nocerino, V.
Colyvas, K. *see* Smart, C.E.
Comlekci, A. *see* Bayraktar, F.
Cooke, D., Hurel, S.J., Casbard, A., Steed, L., Walker, S., Meredith, S., Nunn, A.J., Manca, A., Sculpher, M., Barnard, M., Kerr, D., Weaver, J.U., Ahlquist, J. & Newman, S.P. Randomized controlled trial to assess the impact of continuous glucose monitoring on HbA_{1c} in insulin-treated diabetes (MITRE Study), 540
Cooper, C. *see* Holt, R.I.G.
Cooper, C. *see* Sun, S.
Corbould, A. & Campbell, J. Efficacy of octreotide but not long-acting somatostatin analogue for severe refractory diabetic diarrhoea, 828
Corcoran, P. *see* Collins, M.M.
Cordonnier, M.M. *see* Dumont, I.J.
Corrado, F. *see* D'Anna, R.
Corsi, A. *see* Lepore, G.
Cosson, E., Nguyen, M.T., Pham, I., Pontet, M., Nitenberg, A. & Valensi, P. N-terminal pro-B-type natriuretic peptide: an independent marker for coronary artery disease in asymptomatic diabetic patients, 872
Costa, S. *see* Lepore, G.
Coutinho, J.F. *see* Chaves-Fonseca, R.M.
Cowlam, S. *see* Ripley, D.P.
Craig, D.W., Millis, M.P. & DiStefano, J.K. Genome-wide SNP genotyping study using pooled DNA to identify candidate markers mediating susceptibility to end-stage renal disease attributed to Type 1 diabetes, 1090
Craig, M.E. *see* Catanzariti, L.
Crowne, E. *see* Hambly, H.
Cummings, M. *see* Kar, P.
Cundy, T. *see* McCready, F.
Cuthbertson, J., Patterson, S., O'Harte, F.P.M. & Bell, P.M. Investigation of the effect of oral metformin on dipeptidylpeptidase-4 (DPP-4) activity in Type 2 diabetes, 649
Czupryniak, L. *see* Bialasiewicz, P.

Dabelea, D. *see* Vehik, K.
Dahlén, E.M., Länne, T., Engvall, J., Lindström, T., Grodzinsky, E., Nystrom, F.H. & Östgren, C.J. Carotid intima-media thickness and apolipoprotein B/apolipoprotein A-I ratio in middle-aged patients with Type 2 diabetes, 384
Dahlin, L.B. *see* Thomsen, N.O.B.
Dahlin, L.B. *see* Thomsen, N.O.B.
Dalan, R., Leow, M.K.S., George, J., Chian, K.Y., Tan, A., Han, H.W. & Cheow, S.P. Neuroglycopenia and adrenergic responses to hypoglycaemia: insights from a local epidemic of serendipitous massive overdose of glibenclamide, 105
Dalfrà, M.G. *see* Lapolla, A.
Dalsgaard, E.-M., Lauritzen, T., Christiansen, T., Mai, K.S., Borch-Johnsen, K. & Sandbæk, A. Socioeconomic factors related to attendance at a Type 2 diabetes screening programme, 518
Dalton, R.N. *see* Marcovecchio, M.L.
D'Amato, C. *see* Spallone, V.
Daniel, R.W. *see* Bowling, F.L.
Davidson, D.F. *see* Collier, A.
Davis, T.M.E. *see* Hamilton, E.J.
Davis, T.M.E. *see* Schimke, K.
Davis, W.A. *see* Hamilton, E.J.
Davis, W.A. *see* Schimke, K.
De Cobelli, F. *see* Perseghin, G.
Deakin, T. *see* Cade, J.E.
Debaty, I. *see* Benhamou, P.Y.
Debaty, I. *see* Lablanche, S.
Deedwania, P. *see* Bethel, M.A.
Deguchi, T., Hashiguchi, T., Horinouchi, S., Uto, T., Oku, H., Kimura, K., Makisumi, K. & Arimura, K. Serum VEGF increases

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

in diabetic polyneuropathy, particularly in the neurologically active symptomatic stage, 247
Del Maschio, A. *see* Perseghin, G.
Del Prato, S. Role of glucotoxicity and lipotoxicity in the pathophysiology of Type 2 diabetes mellitus and emerging treatment strategies, 1185
Dellweg, S. *see* Arutchelvam, V.
DeMelo, M. *see* Gucciardi, E.
Demirkan, F. *see* Bayraktar, F.
Dennison, E.M. *see* Holt, R.I.G.
Desmond, D. *see* Coffey, L.
Devemy, F. *see* Dumont, I.J.
Dhatariya, K.K. *see* Sampson, M.J.
Dhatt, G.S. *see* Agarwal, M.M.
Di Benedetto, A. *see* D'Anna, R.
Di Palo, C. *see* Esposito, K.
Diakoumopoulou, E. *see* Tentolouris, N.
DiCianni, G. *see* Lapolla, A.
Diem, P. for the NAVIGATOR Study Group *see* Bethel, M.A.
Diem, P. *see* Krull, I.
Dieterle, C. *see* Kirchberger, I.
DiStefano, J.K. *see* Craig, D.W.
Divers, J. *see* Bleyer, A.J.
Dodesini, A.R. *see* Lepore, G.
Doney, A.S.F., Leese, G.P., Olson, J., Morris, A.D. & Palmer, C.N.A. The Y402H variant of complement factor H is associated with age-related macular degeneration but not with diabetic retinal disease in the Go-DARTS study, 460
Dong, Y.H. *see* Gao, W.G.
Dong, Y.H. *see* Ning, F.
Donnelly, T. *see* Moses, R.G.
Dost, A., Hofer, S., Herbst, A., Stachow, R., Schober, E., Müller, U.A., Holl, R.W. for the Initiative DPV-Science and the German competence network Diabetes. Factors contributing to terminal digital preference in 91 398 patients with diabetes mellitus in Germany and Austria: possible impact on therapeutic decisions, 947
D'Anna, R., Baviera, G., Corrado, F., Giordano, D., Recupero, S. & Di Benedetto, A. First trimester serum neutrophil gelatinase-associated lipocalin in gestational diabetes, 1293
Du, X., Zhu, B., Hu, G., Mao, W., Wang, S., Zhang, H., Wang, F. & Shi, Z. U-shape association between white blood cell count and the risk of diabetes in young Chinese adults, 955
Dukát, A. *see* Price, H.C.
Dumont, I.J., Lepeut, M.S., Tsirtsikolou, D.M., Popielarz, S.M., Cordonnier, M.M., Fayard, A.J., Devemy, F., Fernandez, E., Basuyaux, O. & Jeffcoate, W.J. A proof-of-concept study of the effectiveness of a removable device for offloading in patients with neuropathic ulceration of the foot: the Ransart boot, 778
Dunger, D.B. *see* Marcovecchio, M.L.
Dunseath, G. *see* Peter, R.
Dutta, P. *see* Bansal, D.
Dutta, P. *see* Jayaprakash, P.
Dyce, P. *see* Mohan, K.

Edge, J.A. *see* Smart, C.E.
Edghill, E.L., McCulloch, L., Fulton, P., Beer, N., Hattersley, A.T. & Gloyn, A.L. Mutations in the third gene shown to alter fasting glucose levels in the population (*G6PC2*) are not a common cause of monogenic forms of pancreatic B-cell dysfunction, 113
Edghill, E.L. *see* Rubio-Cabezas, O.
Edmonds, M. *see* Jeffcoate, W.
Eeg-Olofsson, K. *see* Gudbjörnsdottir, S.
Eigenmann, C.A., Colagiuri, R., Skinner, T.C. & Trevena, L. Are current psychometric tools suitable for measuring outcomes of diabetes education?, 425
Eigenmann, C.A. *see* Colagiuri, R.
Ekelund, U. *see* Steele, R.M.
Elbroend, B. *see* Arutchelvam, V.
El-Defrawy, H. *see* Vernon, S.A.
Eliasson, B. *see* Gudbjörnsdottir, S.
Ellard, S. *see* Locke, J.M.
Ellard, S. *see* Shepherd, M.
Ellard, S. *see* Spyer, G.
Ellis, B.M. *see* Paul, L.
Emery, C.J. *see* Choudhary, P.
Émond, V. *see* Ouhoummane, N.
Emslie-Smith, A. *see* Guthrie, B.
Engeland, A. *see* Skurtveit, S.
Englund, E. *see* Thomsen, N.O.B.
Engvall, J. *see* Dahlén, E.M.
Escobedo, J., Buitrón, L.V., Velasco, M.F., Ramírez, J.C., Hernández, R., Macchia, A., Pellegrini, F., Schargrodsky, H., Boissonnet, C. & Champagne, B.M. on behalf of the CARMELA Study Investigators, High prevalence of diabetes and impaired fasting glucose in urban Latin America: the CARMELA Study, 864
Esposito, A. *see* Perseghin, G.
Esposito, K., Maiorino, M.I., Di Palo, C. & Giugliano D. for the Campanian Postprandial Hyperglycemia Study Group, Adherence to a Mediterranean diet and glycaemic control in Type 2 diabetes mellitus, 900
Esteghamati, A., Abbasi, M., Rashidi, A., Meysamie, A., Khalilzadeh, O., Haghazali, M., Asgari, F. & Nakhjavani, M. Optimal waist circumference cut-offs for the diagnosis of metabolic syndrome in Iranian adults: results of the third national survey of risk factors of non-communicable diseases (SuRFNCD-2007), 745
Esteghamati, A., Ashraf, H., Nakhjavani, M., Najafian, B., Hamidi, S. & Abbasi, M. Insulin resistance is an independent correlate of increased urine albumin excretion: a cross-sectional study in Iranian Type 2 diabetic patients, 177
Ewins, D. *see* Ahluwalia, R.

Fadavi, H. *see* Bowling, F.L.
Fagot-Campagna, A. *see* Fosse, S.
Fahrbach, J.L. *see* Wolffenbuttel, B.H.R.
Falvey, H. *see* Hauber, A.B.
Fan, L. *see* Yang, H.
Færch, K., Vaag, A., Witte, D.R., Jørgensen, T., Pedersen, O. & Borch-Johnsen, K. Predictors of future fasting and 2-h post-OGTT plasma glucose levels in middle-aged men and women—the Inter99 Study, 377
Fareedi, M.A. *see* Choudhury, S.M.
Farias, M.G. *see* Chaves-Fonseca, R.M.
Farling, P.A. *see* Silversides, J.A.
Faulks, K. *see* Catanzariti, L.

© 2009 The Authors
Journal compilation © 2009 Diabetes UK.

Fayard, A.J. *see* Dumont, I.J.
Fedele, T. *see* Spallone, V.
Ferland, A., Brassard, P., Lemieux, S., Bergeron, J., Bogaty, P., Bertrand, F., Simard, S. & Poirier, P. Impact of high-fat/low-carbohydrate, high-, low-glycaemic index or low-caloric meals on glucose regulation during aerobic exercise in Type 2 diabetes, 589
Fernandez, E. *see* Dumont, I.J.
Fincher, S. *see* Garg, R.
Finucane, F.M. *see* Steele, R.M.
Fisher, L., Mullan, J.T., Skaff, M.M., Glasgow, R.E., Arean, P. & Hessler, D. Predicting diabetes distress in patients with Type 2 diabetes: a longitudinal study, 622
Fitzgerald, M. *see* Lowe, J.M.
Flack, J. *see* Catanzariti, L.
Flanagan, D.E. The future of glucose sensors—using the technology within a clinical service, 195
Flatman, M. *see* Misra, A.
Fletcher, A.E. *see* Hewitt, J.
Fletcher, J. *see* Burns, J.
Folsom, A.R. *see* Lutsey, P.L.
Forman, E.M. *see* Leeds, J.S.
Fosse, S., Hartemann-Heurtier, A., Jacqueminet, S., Ha Van, G., Grimaldi, A. & Fagot-Campagna, A. Incidence and characteristics of lower limb amputations in people with diabetes, 391
Freckmann, G. *see* Nielsen, J.K.
Freedman, B.I. *see* Bleyer, A.J.
Freeman, D.J. *see* Nelson, S.M.
Fresa, R. *see* Lapolla, A.
Frier, B.M. *see* Major, H.G.
Frontera, G. *see* Rullan, M.
Fujita, H. *see* Narita, T.
Fukuhara, S. *see* Hayashino, Y.
Fulton, P. *see* Edghill, E.L.

Gach, A. *see* Klupa, T.
Gall, M.-A. *see* Moses, R.G.
Gallagher, P. *see* Coffey, L.
Gallen, I.W. *see* Lumb, A.N.
Gama, R. *see* Likhari, T.
Gamble, J. *see* Turzyniecka, M.
Gan, R.M., Wong, V., Cheung, N.W. & McLean, M. Effect of insulin infusion on electrocardiographic findings following acute myocardial infarction: importance of glycaemic control, 174
Gansevoort, R.T. *see* Lambers Heerspink, H.J.
Gao, W.G., Dong, Y.H., Pang, Z.C., Nan, H.R., Zhang, L., Wang, S.J., Ren, J., Ning, F. & Qiao Q. for the Qingdao 2006 Diabetes Survey Group, Increasing trend in the prevalence of Type 2 diabetes and pre-diabetes in the Chinese rural and urban population in Qingdao, China, 1220
Gao, W.G., Qiao, Q., Pitkäniemi, J., Wild, S., Magliano, D., Shaw, J., Söderberg, S., Zimmet, P., Chitson, P., Knowlessur, S., Alberti, G. & Tuomilehto, J. Risk prediction models for the development of diabetes in Mauritian Indians, 996
Gao, W.G. *see* Ning, F.
Gao, X. *see* Yang, H.
García-Morales, E. *see* Aragón-Sánchez, J.
Garg, R., Angus, E. & Fincher, S. Capecitabine-induced severe hypertriglyceridaemia and diabetes: a case report and review of the literature, 1310
Garin, I., Martinez, R., de las Heras, J., Perez-Nanclares, G., Castano, L., Perez de Nanclares, G., the Spanish GEDIMO Group. Mutations in MAFA and IAPP are not a common cause of monogenic diabetes, 746
Garnham, A. *see* Levinger, I.
Gedroyc, W.M.W. *see* Valabhji, J.
George, J. *see* Dalan, R.
Ghezaiel, S. *see* Sun, S.
Ghosh, S. *see* Collier, A.
Giani, G. *see* Rathmann, W.
Giannini, C. *see* Marcovecchio, M.L.
Gianotti, L. *see* Tassone, F.
Gibbs, R.G.J. *see* Valabhji, J.
Gillen, R. *see* Nielsen, J.K.
Giordano, D. *see* D'Anna, R.
Gip, C. *see* Bolin, K.
Giral, P. *see* Béliard, S.
Giugliano, D. for the Campanian Postprandial Hyperglycemia Study Group. *see* Esposito, K.
Glasgow, R.E. *see* Fisher, L.
Gloyn, A.L. *see* Edghill, E.L.
Gloyn, A.L. RD Lawrence Lecture 2009. Old genes, new tricks: learning about blood glucose regulation from naturally occurring genetic variation in humans, 1083
Goenka, N. *see* Ahluwalia, R.
Golay, A. *see* Allet, L.
Gold, A.E. *see* Robertson, H.
Gomm, E.M.W. *see* Patra, S.
Gong, Y. *see* Tong, D.
Goodman, C. *see* Levinger, I.
Gopinath, B. *see* Mitchell, P.
Goran, M.I. *see* Retnakaran, R.
Gosden, C. *see* James, J.
Gosden, C. *see* Nagi, D.
Gosden, C. *see* Nicholson, T.R.J.
Gossain, V.V. *see* Aldasouqi, S.A.
Gough, S.C.L. *see* Manley, S.E.
Gough, S.C.L. *see* Manley, S.E.
Grabert, M. *see* Schober, E.
Grainger, R. *see* Mohan, K.
Greenfield, J.R., Tuthill, A., Soos, M.A., Semple, R.K., Halsall, D.J., Chaudhry, A. & O'Rahilly, S. Severe insulin resistance due to anti-insulin antibodies: response to plasma exchange and immunosuppressive therapy, 79
Greenfield, J.R. *see* Lee, P.
Greenwood, D.C. *see* Cade, J.E.
Greenwood, R.H. *see* Misra, A.
Gregory, J.W. for the DEPICTED Study Team. *see* Hambly, H.
Gregory, J. *see* Allen, D.
Greve, M.D.J. *see* Al-Ansari, S.A.
Griffin, S.J. *see* Steele, R.M.
Grimaldi, A. *see* Fosse, S.
Grodzinsky, E. *see* Dahlén, E.M.
Groenier, K.H. *see* Bilo, H.J.G.
Grossman, E. *see* Weiss, A.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Gucciardi, E., DeMelo, M., Booth, G., Tomlinson, G. & Stewart, D.E. Individual and contextual factors associated with follow-up use of diabetes self-management education programmes: a multisite prospective analysis, 510
Gudbjörnsdottir, S., Eeg-Olofsson, K., Cederholm, J., Zethelius, B., Eliasson, B., Nilsson, P.M. on behalf of the Swedish National Diabetes Register (NDR). Risk factor control in patients with Type 2 diabetes and coronary heart disease: findings from the Swedish National Diabetes Register (NDR), 53
Gupta, R. *see* Agarwal, M.M.
Guthrie, B., Emslie-Smith, A. & Morris, A.D. Which people with Type 2 diabetes achieve good control of intermediate outcomes? Population database study in a UK region, 1269

Ha Van, G. *see* Fosse, S.
Haffner, S.M. *see* Bethel, M.A.
Haghazali, M. *see* Esteghamati, A.
Halsall, D.J. *see* Greenfield, J.R.
Hambly, H., Robling, M., Crowne, E., Hood, K. & Gregory, J.W. for the DEPICTED Study Team. Communication skills of healthcare professionals in paediatric diabetes services, 502
Hamidi, S. *see* Esteghamati, A.
Hamilton, E.J., Rakic, V., Davis, W.A., Chubb, S.A.P., Kamber, N., Prince, R.L. & Davis, T.M.E. Prevalence and predictors of osteopenia and osteoporosis in adults with Type 1 diabetes, 45
Hamilton, J.K. *see* Retnakaran, R.
Hamman, R.F. *see* Vehik, K.
Hammar, N. *see* Ning, F.
Hammond, P. *see* Pickup, J.C.
Han, H.W. *see* Dalan, R.
Han, X.Y. *see* Zhou, X.H.
Hanifi-Moghaddam, P. *see* Kempf, K.
Hanusch-Enserer, U. *see* Stadler, M.
Hardoon, S.L. *see* Thomas, M.C.
Hardy, K. *see* McNulty, S.
Hare, D.L. *see* Levinger, I.
Harper, A. *see* Wong, N.
Harries, L.W. *see* Locke, J.M.
Hartemann-Heurtier, A. *see* Fosse, S.
Haruna, Y. *see* Kawashima, S.
Harvey, E.L. *see* Cade, J.E.
Harvey, J.N. & Lawson, V.L. The importance of health belief models in determining self-care behaviour in diabetes, 5
Hashiguchi, T. *see* Deguchi, T.
Hatazaki, M. *see* Kawashima, S.
Hattersley, A.T. *see* Edghill, E.L.
Hattersley, A.T. *see* Rubio-Cabezas, O.
Hattersley, A.T. *see* Shepherd, M.
Hattersley, A.T. *see* Spyer, G.
Hauber, A.B., Mohamed, A.F., Johnson, F.R. & Falvey, H. Treatment preferences and medication adherence of people with Type 2 diabetes using oral glucose-lowering agents, 416
Hayashino, Y., Fukuhara, S., Akiba, T., Akizawa, T., Asano, Y., Saito, S. & Kurokawa, K. Low health-related quality of life is associated with all-cause mortality in patients with diabetes on haemodialysis: the Japan Dialysis Outcomes and Practice Pattern Study, 921
He, J. *see* Yang, H.
Heidemann, C., Kroll, L., Icks, A., Lampert, T. & Scheidt-Nave, C. Prevalence of known diabetes in German adults aged 25–69 years: results from national health surveys over 15 years, 655
Heier, M. *see* Rathmann, W.
Heinemann, L. *see* Nielsen, J.K.
Heise, T. *see* Arutchelvam, V.
Heise, T. *see* Kapitza, C.
Heller, S.R. *see* Choudhary, P.
de las Heras, J. *see* Garin, I.
Herbst, A. *see* Dost, A.
Hergenç, G. *see* Onat, A.
Hermans, M.P., Ahn, S.A., Selvais, P.L. & Rousseau, M.F. Handedness, insulin sensitivity and pancreatic B-cell function in Type 2 diabetes, 1289
Hernández, R. *see* Escobedo, J.
Hernández-Herrero, M.J. *see* Aragón-Sánchez, J.
Hessler, D. *see* Fisher, L.
Hewitt, J., Smeeth, L., Bulpitt, C.J. & Fletcher, A.E. The prevalence of Type 2 diabetes and its associated health problems in a community-dwelling elderly population, 370
Hierl, F.X. *see* Kirchberger, I.
Hilding, A. *see* Alvarsson, M.
Hill, N.R., Thompson, B., Bruce, J., Matthews, D.R. & Hindmarsh, P. Glycaemic risk assessment in children and young people with Type 1 diabetes mellitus, 740
Hindmarsh, P. *see* Hill, N.R.
Hinz, B.J. *see* Al-Ansari, S.A.
Hire, D. *see* Bleyer, A.J.
Hirota, Y., Ogawa, W., Murawaki, A., Nishiumi, T., Komada, H., Miyake, K., Sakaguchi, K. & Kasuga, M. Deterioration of glycaemic control associated with anti-insulin antibodies likely induced by health supplements, 948
Hofer, S. *see* Dost, A.
Holcombe, J.H. *see* Zhao, Y.
Holl, R.W. for the Initiative DPV-Science. *see* Dost, A.
Holl, R.W. on behalf of the DPV-Wiss Initiative of the German Working Group for Paediatric Diabetology. *see* Schober, E.
Holle, R. *see* Rathmann, W.
Hollins, L. *see* Cade, J.E.
Holt, R.I.G., Phillips, D.I.W., Jameson, K.A., Cooper, C., Dennison, E.M. & Peveler, R.C. the Hertfordshire Cohort Study Group. The relationship between depression and diabetes mellitus: findings from the Hertfordshire Cohort Study, 641
Holt, R.I.G. *see* James, J.
Holt, R.I.G. *see* Nagi, D.
Home, P.D. *see* Arutchelvam, V.
Hood, K. *see* Hambly, H.
Hood, K.K. *see* Williams, L.B.
Hopkins, D. *see* Rogers, H.
Horgan, O. *see* Coffey, L.
Horikawa, A., Ishii-Nozawa, R., Ohguro, M., Takagi, S., Ohtuji, M., Yamada, M., Kuzuya, N., Ujihara, N., Ujihara, M. & Takeuchi, K. Prevalence of GORD (gastro-oesophageal reflux disease) in Type 2 diabetes and a comparison of clinical profiles between diabetic patients with and without GORD, 228
Horinouchi, S. *see* Deguchi, T.
Horvath, K. *see* Siebenhofer, A.
Hosker, J.P. *see* Ndip, A.
Hosoba, M. *see* Narita, T.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Hota, D. *see* Bansal, D.
Hou, J. *see* Xiang, G.D.
Hovorka, R. *see* Brackenridge, A.L.
Hsieh, C.-J. *see* Chen, M.-C.
Hu, G. *see* Du, X.
Hu, Y. *see* Yang, H.
Hugel, G. *see* Johnson, J.A.
Hughes, R. *see* Mohan, K.
Huntsman-Labed, A. *see* Bethel, M.A.
Hurel, S.J. *see* Cooke, D.
Hyer, S.L. *see* Balani, J.

Iafusco, D. *see* Nocerino, V.
Icks, A. *see* Heidemann, C.
Idris, I. *see* Bulugahapitiya, U.
Ilkova, H. *see* Makine, C.
Im, Y.S. *see* Kim, H.C.
Imano, E. *see* Kawashima, S.
Inoue, K., Matsumoto, M. & Akimoto, K. The threshold for definition of impaired fasting glucose in a Japanese population, 1175
Ishii-Nozawa, R. *see* Horikawa, A.
Ishizaki, M. *see* Sakurai, M.
Ismail, K. *see* Nicholson, T.R.J.

Jackson, N. *see* Brackenridge, A.L.
Jacqueminet, S. *see* Fosse, S.
Jaeger, C. *see* Kempf, K.
Jallet, K. *see* Kapitza, C.
James, J., Gosden, C., Winocour, P., Walton, C., Nagi, D., Turner, B., Williams, R. & Holt, R.I.G. Diabetes specialist nurses and role evolvement: a survey by diabetes UK and ABCD of specialist diabetes services 2007, 560
James, J. *see* Nagi, D.
Jameson, K.A. *see* Holt, R.I.G.
Jansen, H., Wijga, A.H., Smit, H.A., Scholtens, S., Kerkhof, M., Koppelman, G.H., de Jongste, J.C. & Stolk, R.P. HbA_{1c} levels in non-diabetic Dutch children aged 8–9 years: the PIAMA birth cohort study, 122
Jayaprakash, P., Bhansali, S., Bhansali, A., Dutta, P. & Anantharaman, R. Magnitude of foot problems in diabetes in the developing world: a study of 1044 patients, 939
Jeffcoate, W., Edmonds, M., Rayman, G., Sherman, C., Stuart, L. & Turner for the Putting Feet First Implementation Group, B. Putting feet first—national guidance at last, 1081
Jeffcoate, W.J. *see* Dumont, I.J.
Jefferson, W. *see* Brackenridge, A.L.
Jeitler, K. *see* Siebenhofer, A.
Jendrike, N. *see* Nielsen, J.K.
Jenkins, A.J. *see* Chen, S.S.H.
Jenkins, C. *see* Misra, A.
Jerums, G. *see* Levinger, I.
Ji, L.N. *see* Zhou, X.H.
Jiang, H. *see* Wolffenbuttel, B.H.R.
Jin, H.Y., Kang, Y.M., Kim, C.Y., Kim, S.H., Liu, W.J., Piao, M.H., Park, J.H., Baek, H.S. & Park, T.S. Morphological comparison of small nerve fibres in gastric mucosa in non-diabetic and Type 2 diabetic subjects, 943
John, U. *see* Schipf, S.
Johnson, F.R. *see* Hauber, A.B.
Johnson, J.A., Balko, S.U., Hugel, G., Low, C. & Svenson, L.W. Increasing incidence and prevalence with limited survival gains among rural Albertans with diabetes: a retrospective cohort study, 1995–2006, 989
Johnston, D.G. *see* Oliver, N.S.
Jones, C. *see* Sampson, M.J.
Jones, C.D. *see* Misra, A.
de Jongste, J.C. *see* Jansen, H.
Joosten, H. *see* Bilo, H.J.G.
Jorgensen, R.S. & Smyth, J.M. Psychosocial distress predicts diabetes: is additional research warranted?, 191
Jørgensen, T. *see* Færch, K.
Jula, A. *see* Koskinen, T.
Juurinen, L., Tiikkainen, M., Saltevo, J., Nikkilä, K., Lanki, H., Leppävuori, E., Kock, T., Teikari-Myyrä, T., Kauppinen-Mäkelin, R., Kotronen, A. & Yki-Järvinen, H. Nateglinide combination therapy with basal insulin and metformin in patients with Type 2 diabetes, 409

Kadıoğlu, P. *see* Makine, C.
Kähönen, M. *see* Koskinen, T.
Kaihura, C. *see* Anderson, J.M.
Kalpen, A. *see* Owings, T.M.
Kalsekar, A. *see* Pratoomsoot, C.
Kamaruddin, N.A. *see* Marre, M.
Kamber, N. *see* Hamilton, E.J.
Kamecke, U. *see* Nielsen, J.K.
Kamoun, M., Charfi, N., Rekik, N., Mnif, M.F., Mnif, F., Kmiha, H., Mnif, Z. & Abid, M. Neurofibromatosis and Type 1 diabetes mellitus: an unusual association, 1180
Kaneko, S. *see* Sakurai, M.
Kaneto, H. *see* Kawashima, S.
Kang, E.S. *see* Park, S.E.
Kang, Y.M. *see* Jin, H.Y.
Kapellen, Th. *see* Schober, E.
Kapitza, C., Heise, T., Birman, P., Jallet, K., Ramis, J. & Balena, R. Pharmacokinetic and pharmacodynamic properties of taspoglutide, a once-weekly, human GLP-1 analogue, after single-dose administration in patients with Type 2 diabetes, 1156
Kapitza, C. *see* Nielsen, J.K.
Kappler, S. *see* Kempf, K.
Kar, P., Laight, D., Rooprai, H.K., Shaw, K.M. & Cummings, M. Effects of grape seed extract in Type 2 diabetic subjects at high cardiovascular risk: a double blind randomized placebo controlled trial examining metabolic markers, vascular tone, inflammation, oxidative stress and insulin sensitivity, 526
Karamitsos, D. *see* Tentolouris, N.
Karanikolas, G. *see* Tentolouris, N.
Karanti, A. *see* Tentolouris, N.
Karşıdağ, K. *see* Makine, C.
Karşıdağ, Ç. *see* Makine, C.
Karvonen, M. *see* Moltchanova, E.V.
Kaskani, E. *see* Tentolouris, N.
Kästenbauer, T. *see* Stadler, M.
Kasuga, M. *see* Hirota, Y.
Katsilambros, N. *see* Tentolouris, N.
Katsuura, Y. *see* Narita, T.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Kauppinen-Mäkelin, R. *see* Juurinen, L.
Kawashima, A. *see* Kawashima, S.
Kawashima, S., Suzuki, M., Kaneto, H., Imano, E., Haruna, Y., Nishimura, Y., Kawashima, A., Hatazaki, M., Matsuoka, T.-A., Yamasaki, Y. & Matsuhisa, M. Insulin resistance and endothelial dysfunction in type 2 diabetic patients with non-alcoholic steatohepatitis, 661
Kaya, H. *see* Onat, A.
Kazakos, K.A. *see* Pikilidou, M.I.
Kellett, J. *see* Price, H.C.
Keltikangas-Järvinen, L. *see* Koskinen, T.
Kempf, K., Manzo, G., Hanifi-Moghaddam, P., Kappler, S., Seissler, J., Jaeger, C., Boehm, B., Roden, M., Kolb, H., Martin, S. & Schloot, N.C. for the PRODIAB Study Group. Effect of combined oral proteases and flavonoid treatment in subjects at risk of Type 1 diabetes, 1311
Kennedy, E.D. *see* Patel, A.
Kerkhof, M. *see* Jansen, H.
Kerr, D. *see* Cooke, D.
Kerr, D. *see* Partridge, H.
Kessler, L. *see* Benhamou, P.Y.
Khalilzadeh, O. *see* Esteghamati, A.
Kido, T. *see* Sakurai, M.
Kikuchi-Taura, A. *see* Makino, H.
Kilpatrick, E.S. *see* Collier, A.
Kim, C.Y. *see* Jin, H.Y.
Kim, D.H. *see* Park, S.E.
Kim, H.C., Cho, Y.J., Ahn, C.W., Park, K.S., Kim, J.C., Nam, J.S., Im, Y.S., Lee, J.E., Lee, S.C. & Lee, H.K. Nerve growth factor and expression of its receptors in patients with diabetic neuropathy, 1228
Kim, J.C. *see* Kim, H.C.
Kim, S.H. *see* Jin, H.Y.
Kim, S.K. *see* Park, S.E.
Kim, T.H. *see* Best, J.H.
Kim, T.N. *see* Yoo, H.J.
Kimura, K. *see* Deguchi, T.
King, B.R. *see* Smart, C.E.
King, L. *see* Bowling, F.L.
Kirchberger, I., Coenen, M., Hierl, F.X., Dieterle, C., Seissler, J., Stucki, G. & Cieza, A. Validation of the international classification of functioning, disability and health (ICF) core set for diabetes mellitus from the patient perspective using focus groups, 700
Kirk, A., Barnett, J., Leese, G. & Mutrie, N. A randomized trial investigating the 12-month changes in physical activity and health outcomes following a physical activity consultation delivered by a person or in written form in Type 2 diabetes: Time2Act, 293
Kirk, S.F.L. *see* Cade, J.E.
Kırmaz, C. *see* Bayraktar, F.
Kishimoto, I. *see* Makino, H.
Klaff, L.J. *see* Wolffenbuttel, B.H.R.
Kleefstra, N. *see* Bilo, H.J.G.
Klein, H.H. *see* Meyer, M.F.
Klingensmith, G.J. *see* Vehik, K.
Klupa, T., Skupien, J., Mirkiewicz-Sieradzka, B., Gach, A., Noczynska, A., Szalecki, M., Kozek, E., Sieradzki, J., Mlynarski, W. & Malecki, M.T. Diabetic retinopathy in permanent neonatal diabetes due to Kir6.2 gene mutations: the results of a minimum 2-year follow-up after the transfer from insulin to sulphonylurea, 663
Kmiha, H. *see* Kamoun, M.
Knowlessur, S. *see* Gao, W.G.
Ko, G. *see* Chan, J.
Kock, T. *see* Juurinen, L.
Koelker, K.H. *see* Nielsen, J.K.
Kokotis, P. *see* Tentolouris, N.
Kolb, H. *see* Kempf, K.
Koliakos, G.G. *see* Pikilidou, M.I.
Komada, H. *see* Hirota, Y.
Kong, A. *see* Chan, J.
Königsrainer, A. *see* Beckert, S.
Kontelis, L. *see* Tentolouris, N.
Koppelman, G.H. *see* Jansen, H.
Kornowski, R. *see* Weiss, A.
Koschinsky, T. *see* Nielsen, J.K.
Koskinen, T., Kähönen, M., Jula, A., Mattsson, N., Laitinen, T., Keltikangas-Järvinen, L., Viikari, J., Välimäki, I., Rönnemaa, T. & Raitakari, O.T. Metabolic syndrome and short-term heart rate variability in young adults. The Cardiovascular Risk in Young Finns Study, 354
Kotronen, A. *see* Juurinen, L.
Kozek, E. *see* Klupa, T.
Krachtis, P. *see* Tentolouris, N.
Krause, R. *see* Schnedl, W.J.
Krentz, A.J. *see* Turzyniecka, M.
Kroll, L. *see* Heidemann, C.
Krull, I., Sahli, R., Diem, P. & Stettler, C. Variability in cross-reactivity of novel insulin analogues in immunometric insulin assays, 1075
Krzyżanowska-Świniarska, B. *see* Ogonowski, J.
Kuczyńska, M. *see* Ogonowski, J.
Kuo, M.-C. *see* Chen, M.-C.
Kurokawa, K. *see* Hayashino, Y.
Kuzuya, N. *see* Horikawa, A.

Lablanche, S., Debaty, I., Renard, E., Bayle, F., Skalli, S. & Benhamou, P.Y. Dramatic improvement of glucose control after plasmapheresis and prednisone in a patient with Type1 diabetes, 951
Laffel, L.M.B. *see* Markowitz, J.T.
Laffel, L.M.B. *see* Williams, L.B.
Laight, D. *see* Kar, P.
Lairon, D. *see* Béliard, S.
Laitinen, T. *see* Koskinen, T.
Lambers Heerspink, H.J., Nauta, F.L., van der Zee, C.P., Brinkman, J.W., Gansevoort, R.T., de Zeeuw, D. & Bakker, S.J.L. Alkalinization of urine samples preserves albumin concentrations during prolonged frozen storage in patients with diabetes mellitus, 556
Lammi, N. *see* Moltchanova, E.V.
Lamoureux, E. *see* Wong, N.
Lampert, T. *see* Heidemann, C.
Lanki, H. *see* Juurinen, L.
Länne, T. *see* Dahlén, E.M.
Lapolla, A., DiCianni, G., Bruttomesso, D., Dalfrà, M.G., Fresa, R., Mello, G., Napoli, A., Romanelli, T., Sciacca, L., Stefanelli,

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

G., Torlone, E. & Mannino, D. Use of insulin detemir in pregnancy: a report on 10 Type 1 diabetic women, 1181
Lasaridis, A.N. *see* Pikilidou, M.I.
Lattuada, G. *see* Perseghin, G.
Lauritzen, T. *see* Dalsgaard, E.-M.
Lawson, V.L. *see* Harvey, J.N.
Lázaro-Martíez, J.L. *see* Aragón-Sánchez, J.
Lázaro-Martínez, J.L. *see* Aragón-Sánchez, J.
Le Coguiec, F. *see* Chan, J.
Le Thi, T.D. *see* Marre, M.
Ledson, M. *see* Mohan, K.
Lee, H.C. *see* Park, S.E.
Lee, H.K. *see* Kim, H.C.
Lee, J.E. *see* Kim, H.C.
Lee, J.H. *see* Park, S.E.
Lee, N. *see* Norris, S.L.
Lee, P., Greenfield, J.R. & Campbell, L.V. Managing young people with Type 1 diabetes in a 'rave' new world: metabolic complications of substance abuse in Type 1 diabetes, 328
Lee, S.C. *see* Kim, H.C.
Leeder, S.R. *see* Mitchell, P.
Leeds, J.S., Forman, E.M., Morley, S., Scott, A.R., Tesfaye, S. & Sanders, D.S. Abnormal liver function tests in patients with Type 1 diabetes mellitus: prevalence, clinical correlations and underlying pathologies, 1235
Leese, G. *see* Kirk, A.
Leese, G.P. *see* Doney, A.S.F.
Leese, G.P. *see* Paul, L.
Leese, G.P. *see* Schofield, C.J.
Lemieux, S. *see* Ferland, A.
Leow, M.K.S. *see* Dalan, R.
Lepeut, M.S. *see* Dumont, I.J.
Lepore, G., Bruttomesso, D., Bonomo, M., Dodesini, A.R., Costa, S., Meneghini, E., Corsi, A., Nosari, I. & Trevisan, R. Continuous subcutaneous insulin infusion is more effective than multiple daily insulin injections in preventing albumin excretion rate increase in Type 1 diabetic patients, 602
Leppävuori, E. *see* Juurinen, L.
Lessa, I. *see* Chaves-Fonseca, R.M.
Levinger, I., Goodman, C., Peake, J., Garnham, A., Hare, D.L., Jerums, G. & Selig, S. Inflammation, hepatic enzymes and resistance training in individuals with metabolic risk factors, 220
Levitt, N.S. *see* Bethel, M.A.
Lezotte, D. *see* Vehik, K.
Li, C.L., Chang, H.Y. & Lu, J.R. Health-related quality of life predicts hospital admission within 1 year in people with diabetes: a nationwide study from Taiwan, 1055
Likhari, T. & Gama, R. Algorithm combining HbA_{1c} and fasting plasma glucose for screening subjects for OGTT, 830
Likhari, T. & Gama, R. Glycaemia-independent ethnic differences in HbA_{1c} in subjects with impaired glucose tolerance, 1068
Lin, F.-Y. *see* Zhang, C.-X.
Lindgren, B. *see* Bolin, K.
Lindow, S.W. *see* Ng, J.M.
Lindsay, R.S. *see* Nelson, S.M.
Lindström, T. *see* Dahlén, E.M.
Ling, G.R., Singh, K., Wong, S.Y. & Toh, V.K.L. Severe neuroglycopaenia secondary to severe hypoglycaemia from serendipitous overdose of gliclazide without adrenergic or autonomic response, 1182
Lipp, R.W. *see* Schnedl, W.J.
Lissett, C.A. *see* Brooks, A.M.S.
Lisy, L. *see* Boaz, M.
Liu, W.J. *see* Jin, H.Y.
Liu, X. *see* Yang, H.
Llobera, J. *see* Rullan, M.
Loba, J. *see* Bialasiewicz, P.
Locke, J.M., Ellard, S., Norwood, V.F. & Harries, L.W. Variants in the isoform-specific coding regions of the HNF1A, HNF4A and HNF1B genes are not a common cause of familial, young-onset diabetes or renal cysts and diabetes (RCAD), 569
Logtenberg, S.J.J. *see* Bilo, H.J.G.
Lonnen, K. *see* Choudhary, P.
Lopes, C. *see* Chaves-Fonseca, R.M.
Lordelo, R.A. *see* Chaves-Fonseca, R.M.
Losekoot, M. *see* Beijers, H.J.B.H.
Low, C. *see* Johnson, J.A.
Lowe, J.M., Mensch, M., McElduff, P., Fitzgerald, M. & Attia, J. Does an advanced insulin education programme improve outcomes and health service use for people with Type 2 diabetes? A 5-year follow-up of the Newcastle Empowerment course, 1277
Lowe, M.R. *see* Markowitz, J.T.
Lu, J.R. *see* Li, C.L.
Lu, Z.X. *see* Manley, S.E.
Lu, Z.X. *see* Manley, S.E.
Lumb, A.N. & Gallen, I.W. Treatment of HNF1-alpha MODY with the DPP-4 inhibitor Sitagliptin, 189
Lunt, H. *see* Moses, R.G.
Luo, Y.Y. *see* Zhou, X.H.
Lutsey, P.L., Pankow, J.S., Bertoni, A.G., Szklo, M. & Folsom, A.R. Serological evidence of infections and Type 2 diabetes: the Multiethnic Study of Atherosclerosis, 149
Luzi, L. *see* Perseghin, G.
Luzio, S.D. *see* Peter, R.
Lyubomirsky, G. *see* Chan, J.

Ma, R. *see* Chan, J.
Macchia, A. *see* Escobedo, J.
Macipe, M. *see* Patra, S.
MacLachlan, M. *see* Coffey, L.
MacLeod, K.M. *see* Choudhary, P.
MacLeod, K.M. *see* Spyer, G.
McCready, F. & Cundy, T. Effects of splenectomy for hereditary spherocytosis on glycated haemoglobin in a woman with Type 2 diabetes, 570
McCulloch, L. *see* Edghill, E.L.
MacDonald, I.A. *see* Choudhary, P.
McElduff, A. *see* Algert, C.S.
McElduff, P. *see* Lowe, J.M.
McEniery, C.M. *see* Anderson, J.M.
McFadyen, A.K. *see* Paul, L.
McLean, M. *see* Gan, R.M.
McMahon, C.M. *see* Mitchell, P.
McMurray, B. *see* Paul, L.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

McNulty, S., Sayed, A. & Hardy, K. Prescription charge exemption and Type 2 diabetes mellitus - a survey of diabetologists' understanding, 827
Magliano, D. *see* Gao, W.G.
Mai, K.S. *see* Dalsgaard, E.-M.
Maiello, M.R. *see* Spallone, V.
Maiorino, M.I. *see* Esposito, K.
Majewski, H. *see* Chen, S.S.H.
Major, H.G., Rees, S.D.R. & Frier, B.M. Driving and diabetes: DVLA response to Lonnen *et al.*, 191
Makine, C., Karşıdağ, Ç., Kadıoğlu, P., Ilkova, H., Karşıdağ, K., Skovlund, S.E., Snoek, F.J. & Pouwer, F. Symptoms of depression and diabetes-specific emotional distress are associated with a negative appraisal of insulin therapy in insulin-naïve patients with Type 2 diabetes mellitus. A study from the European Depression in Diabetes [EDID] Research Consortium, 28
Makino, H., Okada, S., Nagumo, A., Sugisawa, T., Miyamoto, Y., Kishimoto, I., Kikuchi-Taura, A., Soma, T., Taguchi, A. & Yoshimasa, Y. Decreased circulating CD34+ cells are associated with progression of diabetic nephropathy, 171
Makisumi, K. *see* Deguchi, T.
Malecki, M.T. *see* Klupa, T.
Mali, T. *see* Valabhji, J.
Malik, R.A. *see* Thomsen, N.O.B.
Malmberg, K. *see* Ning, F.
Manca, A. *see* Cooke, D.
Manley, S.E., Sikaris, K.A., Lu, Z.X., Nightingale, P.G., Stratton, I.M., Round, R.A., Baskar, V. & Gough, S.C.L. Algorithm combining HbA_{1c} and fasting plasma glucose for screening subjects for OGTT: Authors' response, 831
Manley, S.E., Sikaris, K.A., Lu, Z.X., Nightingale, P.G., Stratton, I.M., Round, R.A., Baskar, V., Gough, S.C.L. & Smith, J.M. Validation of an algorithm combining haemoglobin A_{1c} and fasting plasma glucose for diagnosis of diabetes mellitus in UK and Australian populations, 115
Mannino, D. *see* Lapolla, A.
Manzo, G. *see* Kempf, K.
Mao, W. *see* Du, X.
Maraninchi, M. *see* Béliard, S.
Marcovecchio, M.L., Giannini, C., Dalton, R.N., Widmer, B., Chiarelli, F. & Dunger, D.B. Reduced endogenous secretory receptor for advanced glycation end products (esRAGE) in young people with Type 1 diabetes developing microalbuminuria, 815
Marinou, K. *see* Tentolouris, N.
Markowitz, J.T., Lowe, M.R., Volkening, L.K. & Laffel, L.M.B. Self-reported history of overweight and its relationship to disordered eating in adolescent girls with Type 1 diabetes, 1165
Marre, M., Shaw, J., Brändle, M., Bebakar, W.M.W., Kamaruddin, N.A., Strand, J., Zdravkovic, M., Le Thi, T.D. & Colagiuri, S. on behalf of the LEAD-1 SU study group, Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with Type 2 diabetes (LEAD-1 SU), 268
Martin, S. *see* Kempf, K.
Martin, S. *see* Wolffenbuttel, B.H.R.
Martinez, R. *see* Garin, I.
Martín-Frías, M., Colino, E., Pérez de Nanclares, G., Alonso, M., Ros, P. & Barrio, R. Glibenclamide treatment in relapsed transient neonatal diabetes as a result of a KCNJ11 activating mutation (N48D), 567
Masmiquel, L. *see* Rullan, M.
Masson, E.A. *see* Ng, J.M.
Matos, O.S. *see* Chaves-Fonseca, R.M.
Matsuhisa, M. *see* Kawashima, S.
Matsumoto, M. *see* Inoue, K.
Matsuoka, T.-A. *see* Kawashima, S.
Matteoni-Athayde, L. *see* Chaves-Fonseca, R.M.
Matthews, D. *see* Patel, A.
Matthews, D.J. *see* Bowling, F.L.
Matthews, D.R. *see* Hill, N.R.
Mattsson, N. *see* Koskinen, T.
Mazar, M. *see* Stern, Z.
Mazze, R. *see* Chaves-Fonseca, R.M.
Medlock, R. *see* Ng, J.M.
Meisinger, C. *see* Rathmann, W.
Meller, M.T. *see* Ripley, D.P.
Mello, G. *see* Lapolla, A.
Meneghini, E. *see* Lepore, G.
Mensch, M. *see* Lowe, J.M.
Meredith, S. *see* Cooke, D.
Meyer, M.F., Rose, C.J., Schatz, H. & Klein, H.H. Effects of a short-term improvement in glycaemic control on skin microvascular dysfunction in Type 1 and Type 2 diabetic patients, 880
Meysamie, A. *see* Esteghamati, A.
Miazgowski, T. *see* Ogonowski, J.
Miller, H. *see* Mohan, K.
Milliat-Guittard, L. *see* Benhamou, P.Y.
Millis, M.P. *see* Craig, D.W.
Mindell, J. *see* Pierce, M.B.
Minns, I. *see* Arutchelvam, V.
Mirkiewicz-Sieradzka, B. *see* Klupa, T.
Misra, A., Bachmann, M.O., Greenwood, R.H., Jenkins, C., Shaw, A., Barakat, O., Flatman, M. & Jones, C.D. Trends in yield and effects of screening intervals during 17 years of a large UK community-based diabetic retinopathy screening programme, 1040
Mitchell, J. *see* Brose, L.S.
Mitchell, P., Gopinath, B., McMahon, C.M., Rochtchina, E., Wang, J.J., Boyages, S.C. & Leeder, S.R. Relationship of Type2 diabetes to the prevalence, incidence and progression of age-related hearing loss, 483
Miura, K. *see* Sakurai, M.
Miyake, K. *see* Hirota, Y.
Miyamoto, Y. *see* Makino, H.
Mlynarski, W. *see* Klupa, T.
Mnif, F. *see* Kamoun, M.
Mnif, M.F. *see* Kamoun, M.
Mnif, Z. *see* Kamoun, M.
Mohamed, A.F. *see* Hauber, A.B.
Mohan, K., Miller, H., Dyce, P., Grainger, R., Hughes, R., Vora, J., Ledson, M. & Walshaw, M. Mechanisms of glucose intolerance in cystic fibrosis, 582
Mojaddidi, M. *see* Thomsen, N.O.B.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Molho-Pessach, V. *see* Broshtilova, V.
Moltchanova, E.V., Schreier, N., Lammi, N. & Karvonen, M. Seasonal variation of diagnosis of Type 1 diabetes mellitus in children worldwide, 673
Moon, L. *see* Catanzariti, L.
Morel, P. *see* Benhamou, P.Y.
Morelon, E. *see* Benhamou, P.Y.
Morganti, R. *see* Spallone, V.
Mori, M. *see* Negishi, M.
Morii, T. *see* Narita, T.
Morikawa, Y. *see* Sakurai, M.
Mörk, A.-C. *see* Bolin, K.
Mørland, J. *see* Skurtveit, S.
Morley, S. *see* Leeds, J.S.
Morris, A. *see* Guthrie, B.
Morris, A.D. *see* Doney, A.S.F.
Morris, J.M. *see* Algert, C.S.
Morris, R.W. *see* Thomas, M.C.
Moses, R.G., Bartley, P., Lunt, H., O'Brien, R.C., Donnelly, T., Gall, M.-A., Vesterager, A., Wollmer, P. & Roberts, A. Safety and efficacy of inhaled insulin (AERx® iDMS[1]) compared with subcutaneous insulin therapy in patients with Type 1 diabetes: 1-year data from a randomized, parallel group trial, 260
Mullan, J.T. *see* Fisher, L.
Müller, U.A. *see* Dost, A.
Müller-Godeffroy, E., Treichel, S. & Wagner, V.M. on behalf of the German Working Group for Paediatric Pump Therapy. Investigation of quality of life and family burden issues during insulin pump therapy in children with Type 1 diabetes mellitus—a large-scale multicentre pilot study
Murawaki, A. *see* Hirota, Y.
Mutrie, N. *see* Kirk, A.

Nagi, D., Gosden, C., Walton, C., Winocour, P.H., Turner, B., Williams, R., James, J. & Holt, R.I.G. A national survey of the current state of screening services for diabetic retinopathy: ABCD–Diabetes UK survey of specialist diabetes services 2006, 1303
Nagi, D. *see* James, J.
Nagpal, J., Pande, J.N. & Bhartia, A. A double-blind, randomized, placebo-controlled trial of the short-term effect of vitamin D_3 supplementation on insulin sensitivity in apparently healthy, middle-aged, centrally obese men, 19
Nagumo, A. *see* Makino, H.
Najafian, B. *see* Esteghamati, A.
Nakagawa, H. *see* Sakurai, M.
Nakamura, K. *see* Sakurai, M.
Nakhjavani, M. *see* Esteghamati, A.
Nakhjavani, M. *see* Esteghamati, A.
Nam, J.S. *see* Kim, H.C.
Nan, H.R. *see* Gao, W.G.
Nan, H.R. *see* Ning, F.
Napoli, A. *see* Lapolla, A.
Narita, T., Katsuura, Y., Sato, T., Hosoba, M., Fujita, H., Morii, T. & Yamada, Y. Miglitol induces prolonged and enhanced glucagon-like peptide-1 and reduced gastric inhibitory polypeptide responses after ingestion of a mixed meal in Japanese Type 2 diabetic patients, 187
Naruse, Y. *see* Sakurai, M.
Nauta, F.L. *see* Lambers Heerspink, H.J.
Ndip, A., Basu, A., Hosker, J.P. & Boulton, A.J.M. Diabetic thoracic polyradiculoneuropathy (DTP) following normalization of blood glucose post-pancreatic transplantation, 744
Negishi, M., Shimomura, K., Proks, P., Mori, M. & Shimomura, Y. Mechanism of disopyramide-induced hypoglycaemia in a patient with Type 2 diabetes, 76
Nelson, P. *see* Cade, J.E.
Nelson, S.M., Freeman, D.J., Sattar, N. & Lindsay, R.S. Erythrocytosis in offspring of mothers with Type 1 diabetes—are factors other than insulin critical determinants?, 887
Newman, S.P. *see* Cooke, D.
Ng, J.M., Medlock, R., Wilkinson, L., Lindow, S.W., Masson, E.A. & Allan, B. National Institute for Clinical Excellence (NICE) and the post-partum fasting glucose: are patients with Type 2 diabetes mellitus being missed?, 310
Nguyen, M.T. *see* Cosson, E.
Nicholson, T.R.J., Taylor, J.-P., Gosden, C., Trigwell, P. & Ismail, K. National guidelines for psychological care in diabetes: how mindful have we been?, 447
Nicolaides, K.H. *see* Anderson, J.M.
Nicolay, A. *see* Béliard, S.
Nielsen, J.K., Freckmann, G., Kapitza, C., Ocvirk, G., Koelker, K.H., Kamecke, U., Gillen, R., Amann-Zalan, I., Jendrike, N., Christiansen, J.S., Koschinsky, T. & Heinemann, L. Glucose monitoring by microdialysis: performance in a multicentre study, 714
Nightingale, P.G. *see* Manley, S.E.
Nightingale, P.G. *see* Manley, S.E.
Nikkilä, K. *see* Juurinen, L.
Nikolia, Z. *see* Tentolouris, N.
Nilsson, P.M. *see* Gudbjörnsdottir, S.
Nilsson, P.M. *see* Pikilidou, M.I.
Ning, F., Pang, Z.C., Dong, Y.H., Gao, W.G., Nan, H.R., Wang, S.J., Zhang, L., Ren, J., Tuomilehto, J., Hammar, N., Malmberg, K., Andersson, S.W. & Qiao, Q. for the Qingdao Diabetes Survey Group. Risk factors associated with the dramatic increase in the prevalence of diabetes in the adult Chinese population in Qingdao, China, 855
Ning, F. *see* Gao, W.G.
Nishimura, Y. *see* Kawashima, S.
Nishiumi, T. *see* Hirota, Y.
Nitenberg, A. *see* Cosson, E.
Nocerino, V., Colombo, C., Bonfanti, R., Iafusco, D. & Barbetti, F. Mutation in *IAPP* and *NEUROG3* genes are not a common cause of permanent neonatal/infancy/childhood-onset diabetes, 660
Noczynska, A. *see* Klupa, T.
Nogueira, J.P. *see* Béliard, S.
Norris, J.M. *see* Vehik, K.
Norris, S.L., Lee, N., Thakurta, S. & Chan, B.K.S. Exenatide efficacy and safety: a systematic review, 837
Norwood, V.F. *see* Locke, J.M.
Nosari, I. *see* Lepore, G.
Nowak, D. *see* Bialasiewicz, P.
Ntali, G. *see* Perseghin, G.
Nunn, A.J. *see* Cooke, D.
Nystrom, F.H. *see* Dahlén, E.M.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

O'Rahilly, S. *see* Greenfield, J.R.
O'Brien, R.C. *see* Moses, R.G.
Ocvirk, G. *see* Nielsen, J.K.
Odink, R.J. *see* Beijers, H.J.B.H.
Ogawa, W. *see* Hirota, Y.
Ogonowski, J., Miazgowski, T., Kuczyńska, M., Krzyżanowska-Świniarska, B. & Celewicz, Z. Pregravid body mass index as a predictor of gestational diabetes mellitus, 334
O'Harte, F.P.M. *see* Cuthbertson, J.
Ohguro, M. *see* Horikawa, A.
Ohtuji, M. *see* Horikawa, A.
Okada, S. *see* Makino, H.
Oku, H. *see* Deguchi, T.
Oliveira, M. *see* Chaves-Fonseca, R.M.
Oliver, N. *see* Valabhji, J.
Oliver, N.S., Toumazou, C., Cass, A.E.G. & Johnston, D.G. Glucose sensors: a review of current and emerging technology, 197
Olson, J. *see* Doney, A.S.F.
Onat, A., Hergenç, G., Ayhan, E., Uğur, M., Kaya, H., Tuncer, M. & Can, G. Serum apolipoprotein C-III in high-density lipoprotein: a key diabetogenic risk factor in Turks, 981
Östenson, C.-G. *see* Alvarsson, M.
Östgren, C.J. *see* Dahlén, E.M.
Othman, Y. *see* Agarwal, M.M.
Ouhoummane, N., Abdous, B., Émond, V. & Poirier, P. Impact of diabetes and gender on survival after acute myocardial infarction in the Province of Quebec, Canada—a population-based study, 609
Owens, D.R. *see* Peter, R.
Owings, T.M., Apelqvist, J., Stenström, A., Becker, M., Bus, S.A., Kalpen, A., Ulbrecht, J.S. & Cavanagh, P.R. Plantar pressures in diabetic patients with foot ulcers which have remained healed, 1141

Paisley, A.N., Savage, M.W. & Wiles, P.G. Stabilizing effect of exenatide in a patient with C-peptide-negative diabetes mellitus, 935
Palmer, C.N.A. *see* Doney, A.S.F.
Pande, J.N. *see* Nagpal, J.
Pang, Z.C. *see* Gao, W.G.
Pang, Z.C. *see* Ning, F.
Pankow, J.S. *see* Lutsey, P.L.
Papacosta, A.O. *see* Thomas, M.C.
Park, G.H. *see* Yoo, H.J.
Park, J.H. *see* Jin, H.Y.
Park, K.S. *see* Kim, H.C.
Park, M.S. *see* Yoo, H.J.
Park, S.E., Kang, E.S., Kim, D.H., Kim, S.K., Lee, J.H., Ahn, C.W., Lee, H.C. & Cha, B.S. Effect of *ABCA1* variant on atherogenic dyslipidaemia in patients with Type2 diabetes treated with rosiglitazone, 577
Park, T.S. *see* Jin, H.Y.
Partridge, H. & Kerr, D. Advantages of insulin pump therapy, 1308
Pataky, Z. *see* Allet, L.
Patel, A., Chester, P., Kennedy, E.D. & Matthews, D. An analysis of major UK newspaper articles pertaining to 'diabetes' and 'cure' over 1 year, 452
Paterson, J.A. *see* Bowling, F.L.
Patra, S., Gomm, E.M.W., Macipe, M. & Bailey, C. Interobserver agreement between primary graders and an expert grader in the Bristol and Weston diabetic retinopathy screening programme: a quality assurance audit, 820
Patterson, S. *see* Cuthbertson, J.
Paul, L., Ellis, B.M., Leese, G.P., McFadyen, A.K. & McMurray, B. The effect of a cognitive or motor task on gait parameters of diabetic patients, with and without neuropathy, 234
Pawlowski, M. *see* Bialasiewicz, P.
Peake, J. *see* Levinger, I.
Pearson, D.W.M. *see* Robertson, H.
Pedersen, O. *see* Færch, K.
Pedersen-Bjergaard, U. *see* Akram, K.
Pellegrini, F. *see* Escobedo, J.
Penfornis, A. *see* Benhamou, P.Y.
Perez de Nanclares, G. *see* Garin, I.
Pérez de Nanclares, G. *see* Martín-Frías, M.
Perez-Nanclares, G. *see* Garin, I.
Perkins, B.A. *see* Bril, V.
Perkins, K. *see* Ahluwalia, R.
Perry, I.J. *see* Collins, M.M.
Perseghin, G., Caumo, A., Lattuada, G., De Cobelli, F., Ntali, G., Esposito, A., Belloni, E., Canu, T., Ragogna, F., Scifo, P., Del Maschio, A. & Luzi, L. Elevated fasting plasma C-peptide occurs in non-diabetic individuals with fatty liver, irrespective of insulin resistance, 847
Peter, R., Dunseath, G., Luzio, S.D., Chudleigh, R., Choudhury, S.R. & Owens, D.R. Relative and absolute contributions of postprandial and fasting plasma glucose to daytime hyperglycaemia and HbA_{1c} in subjects with Type 2 diabetes, 974
Peveler, R.C. the Hertfordshire Cohort Study Group, *see* Holt, R.I.G.
Peyrot, M. *see* Best, J.H.
Pham, I. *see* Cosson, E.
Phillips, D.I.W. *see* Holt, R.I.G.
Phillips, P. *see* Schimke, K.
Pia, A. *see* Tassone, F.
Piao, M.H. *see* Jin, H.Y.
Pickup, J.C. & Hammond, P. NICE guidance on continuous subcutaneous insulin infusion 2008: review of the technology appraisal guidance, 1
Pickup, J.C. Author's response, 312
Pieber, T.R. *see* Siebenhofer, A.
Pierce, M.B., Zaninotto, P., Steel, N. & Mindell, J. Undiagnosed diabetes—data from the English longitudinal study of ageing, 679
Pikilidou, M.I., Lasaridis, A.N., Sarafidis, P.A., Befani, C.D., Koliakos, G.G., Tziolas, I.M., Kazakos, K.A., Yovos, J.G. & Nilsson, P.M. Insulin sensitivity increase after calcium supplementation and change in intraplatelet calcium and sodium-hydrogen exchange in hypertensive patients with Type 2 diabetes, 211
Piot, C. *see* Radermecker, R.P.
Piswanger-Soelkner, C. *see* Schnedl, W.J.
Pitkäniemi, J. *see* Gao, W.G.
Poirier, P. *see* Ferland, A.
Poirier, P. *see* Ouhoummane, N.
Polak, B.C.P. *see* Benbassat, J.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Pontet, M. *see* Cosson, E.
Popielarz, S.M. *see* Dumont, I.J.
Portugal, H. *see* Béliard, S.
Pousada, J. *see* Chaves-Fonseca, R.M.
Pouwer, F. *see* Makine, C.
Prager, R. *see* Stadler, M.
Pratoomsoot, C., Smith, H.T., Kalsekar, A., Boye, K.S., Arellano, J. & Valentine, W.J. An estimation of the long-term clinical and economic benefits of insulin lispro in Type 1 diabetes in the UK, 803
Preece, K. *see* Bowling, F.L.
Preiss, D. & Sattar, N. Metabolic syndrome: collapsing under its own weight?, 457
Price, H.C., Thorne, K.I., Dukát, A. & Kellett, J. European physicians overestimate life expectancy and the likely impact of interventions in individuals with Type 2 diabetes, 453
Prince, R.L. *see* Hamilton, E.J.
Procopio, M. *see* Tassone, F.
Proks, P. *see* Negishi, M.
Pu, J.H. *see* Xiang, G.D.
Puttha, R. *see* Sun, S.

Qi, Y. *see* Retnakaran, R.
Qiao, Q. for the Qingdao 2006 Diabetes Survey Group. *see* Gao, W.G.
Qiao, Q. for the Qingdao Diabetes Survey Group. *see* Ning, F.
Qiao, Q. *see* Gao, W.G.
Qiao, Q. *see* Zhou, X.H.
Quehenberger, P. *see* Stadler, M.
Quintana-Marrero, Y. *see* Aragón-Sánchez, J.

Radermecker, R.P., Sultan, A., Piot, C., Remy, A.S., Avignon, A. & Renard, E. Continuous glucose monitoring as a tool to identify hyperglycaemia in non-diabetic patients with acute coronary syndromes, 167
Ragogna, F. *see* Perseghin, G.
Raitakari, O.T. *see* Koskinen, T.
Rakic, V. *see* Hamilton, E.J.
Rami, B. *see* Schober, E.
Ramírez, J.C. *see* Escobedo, J.
Ramis, J. *see* Kapitza, C.
Ramot, Y. *see* Broshtilova, V.
Rashidi, A. *see* Esteghamati, A.
Rathmann, W., Strassburger, K., Heier, M., Holle, R., Thorand, B., Giani, G. & Meisinger, C. Incidence of Type 2 diabetes in the elderly German population and the effect of clinical and lifestyle risk factors: KORA S4/F4 cohort study, 1212
Rayman, G. *see* Jeffcoate, W.
Reaney, M.D. *see* Speight, J.
Recupero, S. *see* D'Anna, R.
Rees, S.D.R. *see* Major, H.G.
Reinehr, Th. *see* Schober, E.
Rekik, N. *see* Kamoun, M.
Remy, A.S. *see* Radermecker, R.P.
Ren, J. *see* Gao, W.G.
Ren, J. *see* Ning, F.
Renard, E. *see* Benhamou, P.Y.
Renard, E. *see* Lablanche, S.
Renard, E. *see* Radermecker, R.P.
Retnakaran, R. & Shah, B.R. Abnormal screening glucose challenge test in pregnancy and future risk of diabetes in young women, 474
Retnakaran, R., Qi, Y., Goran, M.I. & Hamilton, J.K. Evaluation of proposed oral disposition index measures in relation to the actual disposition index, 1198
Ribeiro, M.N. *see* Chaves-Fonseca, R.M.
Ripley, D.P., Wilson, E.J., Meller, M.T. & Cowlam, S. Pneumorrhachis: a rare compliaction of diabetic ketoacidosis, 566
Roberts, A. *see* Moses, R.G.
Roberts, C.L. *see* Algert, C.S.
Robertson, H., Pearson, D.W.M. & Gold, A.E. Severe hypoglycaemia during pregnancy in women with Type 1 diabetes is common and planning pregnancy does not decrease the risk, 824
Robling, M. *see* Hambly, H.
Rochtchina, E. *see* Mitchell, P.
Roden, M. *see* Kempf, K.
Rodin, D.A. *see* Balani, J.
Rogers, H., Turner, E., Thompson, G., Hopkins, D. & Amiel, S.A. Hub-and-spoke model for a 5-day structured patient education programme for people with Type 1 diabetes, 915
Romanelli, T. *see* Lapolla, A.
Rönnemaa, T. *see* Koskinen, T.
Rooprai, H.K. *see* Kar, P.
Ros, P. *see* Martín-Frías, M.
Rose, C.J. *see* Meyer, M.F.
Rosén, I. *see* Thomsen, N.O.B.
Ross, K. *see* Smart, C.E.
Round, R.A. *see* Manley, S.E.
Round, R.A. *see* Manley, S.E.
Rousseau, M.F. *see* Hermans, M.P.
Rubin, R.R. *see* Best, J.H.
Rubio-Cabezas, O., Edghill, E.L., Argente, J. & Hattersley, A.T. Testing for monogenic diabetes among children and adolescents with antibody-negative clinically defined Type 1 diabetes, 1070
Rubio-Cabezas, O. *see* Shepherd, M.
Rullan, M., Cerdà, L., Frontera, G., Masmiquel, L. & Llobera, J. Authors' response to 'Comments on the use of bemiparin in diabetic foot ulcers', 313
Russell, G.B. *see* Bleyer, A.J.
Russell-Jones, D. *see* Brackenridge, A.L.

Sahli, R. *see* Krull, I.
Saito, S. *see* Hayashino, Y.
Sakaguchi, K. *see* Hirota, Y.
Sakurai, M., Miura, K., Takamura, T., Ishizaki, M., Morikawa, Y., Nakamura, K., Yoshita, K., Kido, T., Naruse, Y., Kaneko, S. & Nakagawa, H. J-shaped relationship between waist circumference and subsequent risk for Type 2 diabetes: an 8-year follow-up of relatively lean Japanese individuals, 753
Saltevo, J. *see* Juurinen, L.
Salti, I. on behalf of the Diabetes and Ramadan Study Group. Efficacy and safety of insulin glargine and glimepiride in subjects with Type 2 diabetes before, during and after the period of fasting in Ramadan, 1255
Samarasinghe, D. *see* Valabhji, J.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Sampson, M.J., Singh, H., Dhatariya, K.K., Jones, C., Walden, E. & Bradley, C. Psychometric validation and use of a novel diabetes in-patient treatment satisfaction questionnaire, 729
Sandbæk, A. *see* Dalsgaard, E.-M.
Sanders, D.S. *see* Leeds, J.S.
Sarafidis, P.A. *see* Pikilidou, M.I.
Sato, T. *see* Narita, T.
Sattar, N. *see* Nelson, S.M.
Sattar, N. *see* Preiss, D.
Savage, M.W. *see* Paisley, A.N.
Savvidou, M.D. *see* Anderson, J.M.
Sayed, A. *see* McNulty, S.
Schargrodsky, H. *see* Escobedo, J.
Schatz, H. *see* Meyer, M.F.
Scheen, A., Tan, M.H., Betteridge, D.J., Birkeland, K., Schmitz, O. & Charbonnel, B. on behalf of the PROactive investigators. Long-term glycaemic control with metformin–sulphonylurea–pioglitazone triple therapy in the PROactive Study (PROactive 14), 1033
Scheen, A.J., Tan, M.H., Betteridge, D.J., Birkeland, K., Schmitz, O. & Charbonnel, B. on behalf of the PROactive investigators. Long-term glycaemic effects of pioglitazone compared with placebo as add-on treatment to metformin or sulphonylurea monotherapy in PROactive (PROactive 18), 1242
Scheidt-Nave, C. *see* Heidemann, C.
Scheidt-Nave, C. *see* Schipf, S.
Scherbaum, W.A. *see* Kempf, K.
Schimke, K., Chubb, S.A.P., Davis, W.A., Phillips, P. & Davis, T.M.E. Antiplatelet therapy, *Helicobacter pylori* infection and complicated peptic ulcer disease in diabetes: The Fremantle Diabetes Study, 70
Schipf, S., Schmidt, C.O., Alte, D., Werner, A., Scheidt-Nave, C., John, U., Steveling, A., Wallaschofski, H. & Völzke, H. Smoking prevalence in Type 2 diabetes: results of the Study of Health in Pomerania (SHIP) and the German National Health Interview and Examination Survey (GNHIES), 791
Schloot, N.C. *see* Kempf, K.
Schmidt, C.O. *see* Schipf, S.
Schmitz, O. *see* Bethel, M.A.
Schmitz, O. *see* Scheen, A.
Schmitz, O. *see* Scheen, A.J.
Schnedl, W.J., Piswanger-Soelkner, C., Wallner, S.J., Krause, R. & Lipp, R.W. Agenesis of the dorsal pancreas, 112
Schober, E., Rami, B., Grabert, M., Thon, A., Kapellen, Th., Reinehr, Th., Holl, R.W. on behalf of the DPV-Wiss Initiative of the German Working Group for Paediatric Diabetology and the German Competence Network Diabetes Mellitus. Phenotypical aspects of maturity-onset diabetes of the young (MODY diabetes) in comparison with Type 2 diabetes mellitus (T2DM) in children and adolescents: experience from a large multicentre database, 466
Schober, E. *see* Dost, A.
Schofield, C.J., Yu, N. & Leese, G.P. Decreasing amputation rates in patients with diabetes—a population-based study, 773
Scholtens, S. *see* Jansen, H.
Schreier, N. *see* Moltchanova, E.V.
Sciacca, L. *see* Lapolla, A.
Scifo, P. *see* Perseghin, G.
Scott, A.R. *see* Leeds, J.S.
Sculpher, M. *see* Cooke, D.
Seissler, J. *see* Kempf, K.
Seissler, J. *see* Kirchberger, I.
Selig, S. *see* Levinger, I.
Selvais, P.L. *see* Hermans, M.P.
Semple, R.K. *see* Greenfield, J.R.
Senior, P.A. *see* Al-Ansari, S.A.
Sfikakis, P.P. *see* Tentolouris, N.
Shah, B.R. *see* Retnakaran, R.
Shaw, A. *see* Misra, A.
Shaw, J. *see* Gao, W.G.
Shaw, J. *see* Marre, M.
Shaw, J.E. *see* Cameron, A.J.
Shaw, K.M. *see* Kar, P.
Shehata, H. *see* Balani, J.
Shepherd, M., Shields, B., Ellard, S., Rubio-Cabezas, O. & Hattersley, A.T. A genetic diagnosis of HNF1A diabetes alters treatment and improves glycaemic control in the majority of insulin-treated patients, 437
Shepherd, M. *see* Spyer, G.
Sherman, C. *see* Jeffcoate, W.
Sheu, J.-J. *see* Chen, M.-C.
Shi, Z. *see* Du, X.
Shields, B. *see* Shepherd, M.
Shihabi, Z. *see* Bleyer, A.J.
Shimomura, K. *see* Negishi, M.
Shimomura, Y. *see* Negishi, M.
Shojaee-Moradie, F. *see* Brackenridge, A.L.
Siampli, M. *see* Spallone, V.
Siebenhofer, A., Jeitler, K., Berghold, A., Horvath, K. & Pieber, T.R. Severe hypoglycaemia and glycaemic control in Type 1 diabetes: meta-analysis of multiple daily insulin injections compared with continuous subcutaneous insulin infusion, 311
Sieradzki, J. *see* Klupa, T.
Sikaris, K.A. *see* Manley, S.E.
Sikaris, K.A. *see* Manley, S.E.
Silversides, J.A. & Farling, P.A. Diabetic ketoacidosis guidelines and protection from aspiration pneumonitis, 829
Simard, S. *see* Ferland, A.
Simmons, R.K. *see* Steele, R.M.
Singh, H. *see* Sampson, M.J.
Singh, K. *see* Ling, G.R.
Sithole, J. *see* Bulugahapitiya, U.
Siyambalapitiya, S. *see* Bulugahapitiya, U.
Skae, M. *see* Sun, S.
Skaff, M.M. *see* Fisher, L.
Skalli, S. *see* Lablanche, S.
Skinner, T.C. *see* Eigenmann, C.A.
Skovlund, S.E. *see* Makine, C.
Skrivarhaug, T. *see* Skurtveit, S.
Skupien, J. *see* Klupa, T.
Skurtveit, S., Strøm, H., Skrivarhaug, T., Mørland, J., Bramness, J.G. & Engeland, A. Road traffic accident risk in patients with diabetes mellitus receiving blood glucose-lowering drugs. Prospective follow-up study, 404
Sloggett, A. *see* Thomas, M.C.
Smart, C.E., Ross, K., Edge, J.A., Collins, C.E., Colyvas, K. & King, B.R. Children and adolescents on intensive insulin therapy

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

maintain postprandial glycaemic control without precise carbohydrate counting, 279
Smeeth, L. *see* Hewitt, J.
Smit, H.A. *see* Jansen, H.
Smith, H.T. *see* Pratoomsoot, C.
Smith, J.M. *see* Manley, S.E.
Smyth, J.M. *see* Jorgensen, R.S.
Snoek, F.J. *see* Makine, C.
So, W. *see* Chan, J.
Söderberg, S. *see* Gao, W.G.
Soma, T. *see* Makino, H.
Sommavilla, B. *see* Weitgasser, R.
Soos, M.A. *see* Greenfield, J.R.
Spallone, V., Morganti, R., Siampli, M., Fedele, T., D'Amato, C., Cacciotti, L. & Maiello, M.R. Neuropad as a diagnostic tool for diabetic autonomic and sensorimotor neuropathy, 686
Speight, J., Reaney, M.D. & Barnard, K.D. Not all roads lead to Rome—a review of quality of life measurement in adults with diabetes, 315
Speight, J., Reaney, M.D. & Barnard, K.D. The use of hypothetical scenarios and importance weightings when measuring the impact of diabetes on quality of life. A response to Brose *et al.*, 1077
Spruce, M.C. *see* Bristow, I.R.
Spyer, G., Macleod, K.M., Shepherd, M., Ellard, S. & Hattersley, A.T. Pregnancy outcome in patients with raised blood glucose due to a heterozygous glucokinase gene mutation, 14
Stachow, R. *see* Dost, A.
Stadler, M., Theuer, E., Anderwald, C., Hanusch-Enserer, U., Auinger, M., Bieglmayer, C., Quehenberger, P., Bischof, M., Kästenbauer, T., Wolzt, M., Wagner, O. & Prager, R. Persistent arterial stiffness and endothelial dysfunction following successful pancreas–kidney transplantation in Type 1 diabetes, 1010
Stannard, K. *see* Arun, C.S.
Steed, L. *see* Cooke, D.
Steel, N. *see* Pierce, M.B.
Steele, R.M., Finucane, F.M., Simmons, R.K., Griffin, S.J., Wareham, N.J. & Ekelund, U., on behalf of the ProActive research team. Altered respiratory function is associated with increased metabolic risk, independently of adiposity, fitness and physical activity, 362
Stefanelli, G. *see* Lapolla, A.
Stenström, A. *see* Owings, T.M.
Stern, Z., Calderon-Margalit, R., Mazar, M., Brezis, M. & Tirosh, A. Emergency room visit: a red-flag indicator for poor diabetes care, 1105
Stettler, C. *see* Krull, I.
Steveling, A. *see* Schipf, S.
Stewart, D.E. *see* Gucciardi, E.
Stieglbauer, K. *see* Weitgasser, R.
Stolinski, M. *see* Brackenridge, A.L.
Stolk, R.P. *see* Jansen, H.
Strand, J. *see* Marre, M.
Strassburger, K. *see* Rathmann, W.
Stratton, I.M. *see* Manley, S.E.
Stratton, I.M. *see* Manley, S.E.
Strock, E. *see* Chaves-Fonseca, R.M.
Strøm, H. *see* Skurtveit, S.
Stuart, L. *see* Jeffcoate, W.
Stucki, G. *see* Kirchberger, I.
Sugisawa, T. *see* Makino, H.
Sultan, A. *see* Radermecker, R.P.
Sun, H.L. *see* Xiang, G.D.
Sun, S., Puttha, R., Ghezaiel, S., Skae, M., Cooper, C. & Amin, R. on behalf of the North West England Paediatric Diabetes Network, The effect of biopsy-positive silent coeliac disease and treatment with a gluten-free diet on growth and glycaemic control in children with Type 1 diabetes, 1250
Sundermann, K. *see* Beckert, S.
Suzuki, M. *see* Kawashima, S.
Svenson, L.W. *see* Johnson, J.A.
Swindle, R. *see* Zhao, Y.
Szalecki, M. *see* Klupa, T.
Szklo, M. *see* Lutsey, P.L.

Taguchi, A. *see* Makino, H.
Takagi, S. *see* Horikawa, A.
Takamura, T. *see* Sakurai, M.
Takeuchi, K. *see* Horikawa, A.
Tamesis, B. *see* Chan, J.
Tan, A. *see* Dalan, R.
Tan, M.H. *see* Scheen, A.
Tan, M.H. *see* Scheen, A.J.
Tassone, F., Procopio, M., Gianotti, L., Visconti, G., Pia, A., Terzolo, M. & Borretta, G. Insulin resistance is not coupled with defective insulin secretion in primary hyperparathyroidism, 968
Tavaniotou, E. *see* Tentolouris, N.
Taylor, J.-P. *see* Nicholson, T.R.J.
Taylor, R. *see* Arun, C.S.
Teikari-Myyrä, T. *see* Juurinen, L.
Tennant, M.T.S. *see* Al-Ansari, S.A.
Tentolouris, N., Andrianakos, A., Karanikolas, G., Karamitsos, D., Trontzas, P., Krachtis, P., Christoyannis, F., Tavaniotou, E., Nikolia, Z., Kaskani, E., Kontelis, L. & Sfikakis, P.P. Prevalence of diabetes mellitus and obesity in the general adult population of Greece: a door-to-door epidemiological study, 451
Tentolouris, N., Marinou, K., Kokotis, P., Karanti, A., Diakoumopoulou, E. & Katsilambros, N. Sudomotor dysfunction is associated with foot ulceration in diabetes, 302
Terzolo, M. *see* Tassone, F.
Tesfaye, S. *see* Leeds, J.S.
Thakurta, S. *see* Norris, S.L.
Theuer, E. *see* Stadler, M.
Thivolet, C. *see* Benhamou, P.Y.
Thomas, M.C., Hardoon, S.L., Papacosta, A.O., Morris, R.W., Wannamethee, S.G., Sloggett, A. & Whincup, P.H. Evidence of an accelerating increase in prevalence of diagnosed Type 2 diabetes in British men, 1978–2005, 766
Thompson, B. *see* Hill, N.R.
Thompson, G. *see* Rogers, H.
Thomsen, N.O.B., Englund, E., Thrainsdottir, S., Rosén, I. & Dahlin, L.B. Intraepidermal nerve fibre density at wrist level in diabetic and non-diabetic patients, 1120
Thomsen, N.O.B., Mojaddidi, M., Malik, R.A. & Dahlin, L.B. Biopsy of the posterior interosseous nerve: a low morbidity method for assessment of peripheral nerve disorders, 100

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Thon, A. *see* Schober, E.
Thorand, B. *see* Rathmann, W.
Thorne, K.I. *see* Price, H.C.
Thorsteinsson, B. *see* Akram, K.
Thrainsdottir, S. *see* Thomsen, N.O.B.
Tiikkainen, M. *see* Juurinen, L.
Tikellis, G. *see* Wong, N.
Tirosh, A. *see* Stern, Z.
Toh, V.K.L. *see* Ling, G.R.
Tomioka, S. *see* Bril, V.
Tomlinson, G. *see* Gucciardi, E.
Tong, D. & Gong, Y. What causes irreversible prolonged coma after severe hypoglycaemia?, 749
Tong, P. *see* Chan, J.
Torlone, E. *see* Lapolla, A.
Toso, C. *see* Benhamou, P.Y.
Toumazou, C. *see* Oliver, N.S.
Treichel, S. *see* Müller-Godeffroy, E.
Trevena, L. *see* Eigenmann, C.A.
Trevisan, R. *see* Lepore, G.
Trigwell, P. *see* Nicholson, T.R.J.
Trontzas, P. *see* Tentolouris, N.
Tse, L.-A. *see* Zhang, C.-X.
Tsirtsikolou, D.M. *see* Dumont, I.J.
Tuncer, M. *see* Onat, A.
Tuomilehto, J. *see* Gao, W.G.
Tuomilehto, J. *see* Ning, F.
Turner for the Putting Feet First Implementation Group, B. *see* Jeffcoate, W.
Turner, B. *see* James, J.
Turner, B. *see* Nagi, D.
Turner, E. *see* Rogers, H.
Turzyniecka, M., Wild, S.H., Krentz, A.J., Chipperfield, A.J., Gamble, J., Clough, G.F. & Byrne, C.D. Skeletal muscle microvascular exchange capacity is associated with hyperglycaemia in subjects with central obesity, 1112
Tuthill, A. *see* Greenfield, J.R.
Tziolas, I.M. *see* Pikilidou, M.I.

Ubink-Veltmaat, L.J. *see* Bilo, H.J.G.
Uğur, M. *see* Onat, A.
Ujihara, M. *see* Horikawa, A.
Ujihara, N. *see* Horikawa, A.
Ulbrecht, J.S. *see* Owings, T.M.
Umpleby, A.M. *see* Brackenridge, A.L.
Uto, T. *see* Deguchi, T.

Vaag, A. *see* Færch, K.
Valabhji, J., Oliver, N., Samarasinghe, D., Mali, T., Gibbs, R.G.J. & Gedroyc, W.M.W. Conservative management of diabetic forefoot ulceration complicated by underlying osteomyelitis: the benefits of magnetic resonance imaging, 1127
Valensi, P. *see* Cosson, E.
Valentine, W.J. *see* Pratoomsoot, C.
Valéro, R. *see* Béliard, S.
Välimäki, I. *see* Koskinen, T.
Vehik, K., Hamman, R.F., Lezotte, D., Norris, J.M., Klingensmith, G.J. & Dabelea, D. Childhood growth and age at diagnosis with Type 1 diabetes in Colorado young people, 961
Velasco, M.F. *see* Escobedo, J.
Vernon, S.A., Bhagey, J., Boraik, M. & El-Defrawy, H. Long-term review of driving potential following bilateral panretinal photocoagulation for proliferative diabetic retinopathy, 97
Vesterager, A. *see* Moses, R.G.
Vialettes, B. *see* Béliard, S.
Vicaretti, M. *see* Burns, J.
Viikari, J. *see* Koskinen, T.
Visconti, G. *see* Tassone, F.
Viswanathan, G.N. Diabetic Cardiology—Fisher M, McMurray J eds. *reviewed by*, 751
Volkening, L.K. *see* Markowitz, J.T.
Völzke, H. *see* Schipf, S.
Vora, J. *see* Mohan, K.

Wagner, V.M. on behalf of the German Working Group for Paediatric Pump Therapy, V.M. *see* Müller-Godeffroy, E.
Wagner, O. *see* Stadler, M.
Wainstein, J. *see* Boaz, M.
Walden, E. *see* Sampson, M.J.
Walker, S. *see* Cooke, D.
Wallaschofski, H. *see* Schipf, S.
Wallner, S.J. *see* Schnedl, W.J.
Walshaw, M. *see* Mohan, K.
Walton, C. *see* James, J.
Walton, C. *see* Nagi, D.
Wang, F. *see* Du, X.
Wang, J.J. *see* Mitchell, P.
Wang, J.J. *see* Wong, N.
Wang, P.-W. *see* Chen, M.-C.
Wang, S. *see* Du, X.
Wang, S.J. *see* Gao, W.G.
Wang, S.J. *see* Ning, F.
Wang, S.S. *see* Wong, N.
Wannamethee, S.G. *see* Thomas, M.C.
Wareham, N.J. *see* Steele, R.M.
Waters, A.-M. *see* Catanzariti, L.
Weaver, J.U. *see* Cooke, D.
Wegener, C. *see* Burns, J.
Wei, Y. *see* Yang, H.
Weiss, A., Boaz, M., Beloosesky, Y., Kornowski, R. & Grossman, E. Body mass index and risk of all-cause and cardiovascular mortality in hospitalized elderly patients with diabetes mellitus, 253
Weitgasser, R., Sommavilla, B. & Stieglbauer, K. Insulin treatment in a patient with Type 2 diabetes and Stiff-person syndrome, 111
Werner, A. *see* Schipf, S.
Whincup, P.H. *see* Thomas, M.C.
Widmer, B. *see* Marcovecchio, M.L.
Wijga, A.H. *see* Jansen, H.
Wild, S. *see* Gao, W.G.
Wild, S.H. *see* Turzyniecka, M.
Wiles, P.G. *see* Paisley, A.N.
Wilkinson, L. *see* Ng, J.M.
Williams, L.B., Laffel, L.M.B. & Hood, K.K. Diabetes-specific family conflict and psychological distress in paediatric Type 1 diabetes, 908
Williams, R. *see* Choudhury, S.M.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

Williams, R. *see* Choudhury, S.M.
Williams, R. *see* James, J.
Williams, R. *see* Nagi, D.
Wilson, E.J. *see* Ripley, D.P.
Winocour, P. *see* James, J.
Winocour, P.H. *see* Nagi, D.
Witte, D.R. *see* Færch, K.
Wojtusciszyn, A. *see* Benhamou, P.Y.
Wolf, S. *see* Beckert, S.
Wolffenbuttel, B.H.R., Klaff, L.J., Bhushan, R., Fahrbach, J.L., Jiang, H. & Martin, S. Initiating insulin therapy in elderly patients with Type 2 diabetes: efficacy and safety of lispro mix 25 vs. basal insulin combined with oral glucose-lowering agents, 1147
Wollmer, P. *see* Moses, R.G.
Wolthers, T. *see* Chan, J.
Wolzt, M. *see* Stadler, M.
Wong, N., Wang, S.S., Lamoureux, E., Wong, T.Y., Tikellis, G., Harper, A. & Wang, J.J. Blood pressure control and awareness among patients with diabetes and hypertension attending a tertiary ophthalmic clinic, 34
Wong, R. *see* Chan, J.
Wong, S.Y. *see* Ling, G.R.
Wong, T.Y. *see* Wong, N.
Wong, V. *see* Gan, R.M.

Xiang, G.D., Pu, J.H., Zhao, L.S., Sun, H.L., Hou, J. & Yue, L. Association between plasma osteoprotegerin concentrations and urinary albumin excretion in Type 2 diabetes, 397
Xu, J. *see* Bleyer, A.J.
Xu, X. *see* Yang, H.

Yamada, M. *see* Horikawa, A.
Yamada, Y. *see* Narita, T.
Yamamoto-Suganuma, R. & Aso, Y. Relationship between post-occlusive forearm skin reactive hyperaemia and vascular disease in patients with Type 2 diabetes—a novel index for detecting micro- and macrovascular dysfunction using laser Doppler flowmetry, 83
Yamasaki, Y. *see* Kawashima, S.
Yang, H., Wei, Y., Gao, X., Xu, X., Fan, L., He, J., Hu, Y., Liu, X., Chen, X., Yang, Z. & Zhang, C. on behalf of China National GDM Survey Working Group, Risk factors for gestational diabetes mellitus in Chinese women—a prospective study of 16 286 pregnant women in China, 1099
Yang, S.J. *see* Yoo, H.J.
Yang, X. *see* Chan, J.
Yang, Z. *see* Yang, H.
Ye, W. *see* Zhao, Y.
Ye, X.-Q. *see* Zhang, C.-X.
Yener, S. *see* Bayraktar, F.
Yesil, S. *see* Bayraktar, F.
Yki-Järvinen, H. *see* Juurinen, L.
Yoo, H.J., Park, M.S., Kim, T.N., Yang, S.J., Cho, G.J., Baik, S.H., Choi, D.S., Park, G.H. & Choi, K.M. A Ubiquitous Chronic Disease Care system using cellular phones and the internet, 628
Yoshimasa, Y. *see* Makino, H.
Yoshita, K. *see* Sakurai, M.
Yovos, J.G. *see* Pikilidou, M.I.
Yu, N. *see* Schofield, C.J.
Yue, L. *see* Xiang, G.D.

Zaman, B. *see* Choudhury, S.M.
Zandman-Goddard, G. *see* Boaz, M.
Zaninotto, P. *see* Pierce, M.B.
Zdravkovic, M. *see* Marre, M.
van der Zee, C.P. *see* Lambers Heerspink, H.J.
de Zeeuw, D. *see* Lambers Heerspink, H.J.
Zethelius, B. *see* Gudbjörnsdottir, S.
Zhang, C. on behalf of China National GDM Survey Working Group. *see* Yang, H.
Zhang, C.-X., Tse, L.-A., Ye, X.-Q., Lin, F.-Y., Chen, Y.-M. & Chen, W.-Q. Moderating effects of coping styles on anxiety and depressive symptoms caused by psychological stress in Chinese patients with Type 2 diabetes, 1287
Zhang, H. *see* Du, X.
Zhang, L. *see* Gao, W.G.
Zhang, L. *see* Ning, F.
Zhang, X.Y. *see* Zhou, X.H.
Zhao, L.S. *see* Xiang, G.D.
Zhao, Y., Ye, W., Boye, K.S., Holcombe, J.H. & Swindle, R. Healthcare charges and utilization associated with diabetic neuropathy: impact of Type 1 diabetes and presence of other diabetes-related complications and comorbidities, 61
Zhou, X.H., Ji, L.N., Luo, Y.Y., Zhang, X.Y., Han, X.Y. & Qiao, Q. Performance of HbA_{1c} for detecting newly diagnosed diabetes and pre-diabetes in Chinese communities living in Beijing, 1262
Zhu, B. *see* Du, X.
Zimmet, P. *see* Gao, W.G.
Zimmet, P.Z. *see* Cameron, A.J.
Zlotogorski, A. *see* Broshtilova, V.

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

DOI: 10.1111/j.1464-5491.2009.02884.x

Subject Index

abdominal obesity, 864
accuracy, 714
acute coronary syndrome, 167
adherence, 416
adhesion molecules, 1010
adolescence, 1165
adolescents, 466
aetiology, 377, 1235
African-American, 128
age at diagnosis, 961
age-related macular degeneration, 460
alanine aminotransferase, 1204
albuminuria, 171, 556
alkalinization, 556
All-or-nothing quality measures, 1269
amitriptyline, 1019
amputation, 773, 778, 1063
amputations, 391
anthropometry, 377
anti-insulin antibodies, 79
anxiety, 153, 908, 1287
apolipoprotein C-III, 981
apolipoproteins, 384
apoprotein C-II, 736
apoprotein C-III, 736
Arterial stiness, 1135
Asia, 693
Asian diabetes, 177
aspart insulin, 260
aspirin, 708
associated health problems, 370
atherogenic dyslipidaemia, 577
ATP-binding cassette transporter 1, 577
Australia, 596
autonomic nervous system, 354
autonomic neuropathy, 686
autonomic response, 105
awareness, 34

Bangladeshi, 40
barefoot pressure measurement, 1141
basal/bolus, 260
B-cell function, 1185, 1198
B-cell mass, 1185
Bengladeshi, 636
beta cell function, 345
Birthweight, 1193
blindness, 489
blood pressure control, 34
Blue Mountains Eye Study, 483
Blue Mountains Hearing Study, 483
body image, 1063
body mass index, 253, 334
bone mineral density, 45
bone turnover, 45
Brazil, 286

Caesarean section, 1193
calcium supplementation, 211
capillary blood cell velocity, 880
cardiac autonomic neuropathy, 302
cardiovascular disease, 83, 253, 384, 457
cardiovascular disease risk factors, 1269
cardiovascular risk factors, 864
Carnosinase 1, 1090
carotid intima-media thickness, 864
carpal tunnel syndrome, 1120
Caucasian, 128
caucasians, 1068
cellulitis, 548
cgms, 167
child, 596
children, 122, 466, 493, 673
China, 1099
Chinese, 1220
Chinese adults, 955
chronic disease, 628
chronic disease self-management education, 510
cigarette smoking, 981
clinical aspects, 466
clinical care, 560
clinical score, 240
closed loop, 197
cocaine, 328
coeliac disease, 1250
cognitive function, 665
combination therapy, 1242
communication skills, 502
complement factor H, 460
compliance, 518
complications, 228, 773, 928, 1301
complications of diabetes, 391
Comprehensive ICF Core Set, 700
consultation–liaison psychiatry, 447
Continuous glucose monitoring, 197, 339, 540, 714
continuous subcutaneous insulin infusion, 279
coping, 1287
corneal nerve fibre, 1228
coronary artery disease, 872
coronary heart disease, 53, 981
coronary risk equivalent, 142
cost, 928
cost-eectiveness analysis, 803
C-peptide negative, 935
cross-sectional studies, 655
CSII, 493, 602

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

cyclooxygenase, 182
cystic fibrosis, 582
cystic fibrosis-related diabetes, 582
cytokines, 220
depression, 28, 153, 641, 908, 1287
diabetes, 5, 40, 79, 97, 100, 134, 142, 149, 153, 234, 247, 253, 286, 315, 370, 391, 416, 447, 457, 474, 493, 502, 622, 628, 636, 641, 686, 693, 729, 736, 837, 921, 928, 955, 989, 1019, 1055, 1063, 1105, 1120, 1262, 1277, 1294
diabetes education, 425
diabetes glycaemic control, 740
diabetes management, 34
diabetes mellitus, 28, 53, 128, 609, 655, 700, 935, 939
diabetes patient education, 442
diabetes specialist nurses, 560
diabetes Type 2, 981
diabetes-related complication or comorbidity, 61
diabetic foot, 93, 548, 552, 773, 778, 939, 1127
diabetic foot ulcers, 89
diabetic neuropathy, 61, 240, 1228
diabetic patient, 404
diabetic retinopathy, 34, 460, 489, 783, 1040
diabetic screening, 820
diagnosis, 240, 641, 679, 686, 1175
diagnosis of diabetes mellitus, 115
diet, 900, 1048
dipeptidyl peptidase-4, 268
dipeptidylpeptidase-4, 649
discrete choice experiment, 416
disopyramide, 76
disposition index, 1198
distress, 622, 908
diurnal hyperglycaemia, 974
driving, 97
dual-task interference, 234

echocardiography, 872
economics, 928
education, 915, 1282
elderly, 253, 370, 1147
emergency ward, 1105
empowerment, 1282
endogenous secretory receptor for advanced glycation end products, 815
endothelial dysfunction, 397
endothelium, 171
end-stage renal disease, 1090
environment, 673
epidemiology, 362, 391, 483, 596, 641, 673, 989, 1175
erysipelas, 548
ethnic dierences, 1068
ethnic minority, 636
exenatide, 722, 837, 935
exercise, 589, 1027
expert patient, 1048

falls, 234
family management, 908
fasting glucose, 760
fasting plasma glucose, 115, 974
FEV_1, 362
flavonoids, 526
foot, 893
foot at risk, 939
foot infection, 552
foot ulceration, 302, 939
free fatty acids, 847
frozen storage, 556
FVC, 362

gait, 234
gastric mucosa, 943
gastro-oesophageal reflex disease, 228
gender, 609
general practice, 518
genetic association, 577
genetics, 1083
genome-wide association study, 1090
German National Health Interview and Examination Survey 98, 791
gestational diabetes, 14, 334, 760, 798, 1099, 1298
glargine, 1147
glibenclamide, 105
glomerular filtration rate, 478
glucagon-like peptide-1, 649
glucagon-like peptide-1 receptor agonist, 268
glucagon-like peptide 1 analogue, 1156
glucokinase, 14
glucometer, 760
glucoregulation, 589
glucose, 128, 409
glucose challenge test, 474
glucose intolerance, 474
Glucose sensors, 197
glucotoxicity, 1185
gluten-free diet, 1250
glycaemia in paediatrics, 740
glycaemic assessment, 740
glycaemic control, 837, 880, 900, 1250
glycaemic index, 589
Glycaemic Risk Assessment in Diabetes Equation, 740
glycated haemoglobin, 115, 122, 128, 286, 409, 1112, 1262
glycemic control, 1301
GoKinD study, 1090
grader, 820
grape seed extract, 526

haematocrit, 887
haemodialysis, 89, 921
handedness, 1294
HbA_{1c}, 34, 900, 974, 1068, 1112
healing, 778
health beliefs, 636
health care delivery, 1301
health service utilization, 510
health services, 447
health surveys, 655

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

healthcare charges, 61
healthcare delivery, 679
health-care professionals, 502
healthcare quality, 1269
health-related quality of life, 617, 921, 1055
hearing loss, 483
heart rate variability, 354
height, 961
Helicobacter pylori, 70
hepatic disease, 1235
hepatocyte nuclear factor 1A, 1172
hepatocyte nuclear factor 4A, 1172
high-density lipoprotein, 981
histone acetylation, 182
HNF1A, 437
HOMA, 345
homeostasis model assessment, 847
hospital admission, 1055
hyperglycaemia, 167, 174
hyperinsulinaemic clamp, 211
hyperlipidaemia, 53
hypertension, 53, 628
hypertriglyceridaemia, 736
hypoglycaemia, 76, 105, 409, 665

idiopathic diabetes, 1070
illness cognitions, 5
imaging, 93
impaired fasting glucose, 345, 864, 1212
impaired glucose tolerance, 345, 582, 1068, 1204, 1212
incidence, 596, 753, 955, 1212
incretin, 268
Indian, 996
indicators, 425, 442
infection, 149
inflammation, 182, 526, 708
inhaled insulin, 260
in-patient, 729
INS, 1070
in-shoe pressure measurement, 1141
insulin, 28, 404, 887, 935, 1282
insulin analogues, 279, 1147
insulin detemir, 1027
insulin dose, 409
insulin gene, 1070
insulin glargine, 1027, 1255
insulin infusion, 174
insulin lispro, 803
insulin resistance, 79, 177, 345, 362, 582, 753
insulin secretion, 345, 582, 753, 968, 1185
insulin sensitivity, 19, 211, 968, 1112, 1294
insulin therapy, 602, 915
insulinotropic, 268
intensive care, 167
intermediate density lipoprotein, 532
International Classification of Functioning, 700
internet, 628
interobserver agreement, 820
intervention, 293
intra-epidermal nerve fibre density, 1120
intra-hepatic fat, 847

K_{ATP} channel, 76
ketamine, 328
ketoacidosis, 328

laser Doppler anemometry, 880
laser Doppler flowmetry, 83
laser photocoagulation, 97
laterality, 1294
leg ischaemia, 134
length of stay, 729
leptin, 887
lifestyle, 293, 377
lipid metabolism, 384
lipocalin, 1298
lipoprotein lipase, 736
lipotoxicity, 1185
lispro, 1147
liver enzymes, 220
longitudinal study, 955
low-density lipoprotein, 532

macrovascular function, 83
maculopathy, 489, 1040
magnetic resonance imaging, 1127
magnetic resonance spectroscopy, 847
management, 693
maturity-onset diabetes of the young, 437, 1083, 1172
measurement, 93, 315
measurement tools, 425
meta-analysis, 142
metabolic risk factors, 220
metabolic syndrome, 306, 354, 362, 457, 1204
metformin, 649, 798, 1033, 1242
methicillin-resistant *Staphylococcus aureus*, 552
microalbuminuria, 177, 602, 815
microangiopathy, 171
microcirculation, 89, 1112
microdialysis, 714
microvascular complications, 1090
microvascular dysfunction, 880
microvascular function, 83
mix 25, 1147
models of care, 693
modified Toronto Clinical Neuropathy Score, 240
MODY, 466
monogenic diabetes, 437, 1070
mortality, 253, 478, 921, 989
mycophenolate, 79
myocardial infarction, 174, 609

national consensus, 442
neonatal diabetes, 1083
nerve biopsy, 100
nerve fibre, 943
nerve growth factor, 1228
nerve growth factor receptor, 1228

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

neuroglycopenia, 105
neuropathy, 100, 247, 773, 943, 1003, 1019, 1141
neuropathy disability score, 1228
neutrophil gelatinase-associated lipocalin, 1298
non-alcoholic fatty liver disease, 1204
non-diabetic, 122
non-esterified fatty acid, 532
non-nutritive sweeteners, 589
Norway, 404
NPH insulin, 1027
N-terminal pro-B-type natriuretic peptide, 872

obese, 19
obesity, 306, 362, 628, 855, 1112, 1165
onychomycosis, 548
ooading, 778
oral agents, 416
oral glucose tolerance test, 115, 377, 1198
oral glucose-lowering agent, 404
orthotic, 893
osteomyelitis, 552, 1127
osteopenia, 45
osteoporosis, 45
osteoprotegerin, 397
outcomes, 425, 442, 693
overload hypothesis, 961
oxidative stress, 526

paediatrics, 279, 502
pain, 893, 1019
pancreas transplantation, 1010
pancreatic islet transplantation, 617
participation, 518
pathogen, 149
patient education, 5, 510
patient perspective, 700
patient preferences, 416
peer education, 40
peptic ulcer disease, 70
peripheral neuropathy, 234, 302, 686, 939
peroxisome proliferators-activated receptor-γ agonist, 532
pharmacodynamics, 1156
pharmacogenetics, 1083
pharmacokinetics, 1156
pharmacotherapy, 228
physical activity, 293
physical inactivity, 855
pioglitazone, 1033, 1242
plasma exchange, 79
platelets, 887
polysomnography, 339
population studies, 641
posterior interosseous nerve, 100
postprandial glucose, 279, 900, 974
pre-diabetes, 1220
pregabalin, 1019
pregnancy, 474, 798, 824, 1099, 1135, 1193
pregnancy planning, 824
pressure, 893
prevalence, 370, 655, 766, 989, 1220
prevention, 1003
primary care, 286
primary hyperparathyroidism, 968
productivity, 928
prognosis, 693
proliferative retinopathy, 489
proportional hazards models, 478
prostaglandins, 182
protein gene product 9.5, 1120
psychological stress, 1287
psychology, 447
psychosocial, 5
psychosocial adjustment, 1063
public health, 791
pulse wave analysis, 1135
pulse-wave velocity, 1010

QT interval, 174
qualitative, 636
qualitative method, 700
quality, 1105
quality of life, 315, 493, 893
QUEST questionnaire, 228
questionnaire, 729

Ramadan, 1255
randomized controlled trial, 1048
reactive hyperaemia, 83
record linkage, 1193
registry, 693
regular human insulin, 803
reliability, 783
REM sleep, 339
reproducibility of results, 783
retinal screening, 489
retinopathy, 1277, 1301
review, 315
risk, 457, 1040
risk assessment, 478, 1055
risk factor, 1099
risk factors, 377
risk prediction, 306, 693
risk prediction model, 996
role development, 560
rosiglitazone, 577
rural, 989

safety, 1156
satisfaction, 729
screening, 115, 518, 679, 760, 783, 1040, 1262, 1277
secondary prevention, 53
secular trends, 1220
serological testing, 70
seropositivity, 149
service fragmentation, 560
service provision, 162, 915
severe hypoglycaemia, 824
silent myocardial ischaemia, 872

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.

skin biopsy, 1120
smoking, 1212
smoking prevalence, 791
socioeconomic factors, 518
socio-economic status, 855
South Asians, 1068
staged diabetes management, 286
statins, 708
stem cells, 134
Study of Health in Pomerania, 791
substance abuse, 328
sudomotor function, 302
sulphonylurea, 1033, 1242
sulphonylurea sensitivity, 437
sulphonylureas, 76, 665
sural nerve, 100
survey, 447
survival, 609
sympathetic skin response, 302
symptoms, 665
systematic review, 837

Taiwan, 1055
taspoglutide, 1156
technology, 197
teenagers, 162
temporal trends, 961
testing, 1003
therapy, 1003
thiazolidinedione, 268
threshold, 1141
time trends, 766
tinea pedis, 548
Toronto Clinical Neuropathy Score, 240
total contact cast, 778
trac accident risk, 404
training needs, 502
treatment satisfaction, 722
trend, 596, 655
triple therapy, 1033
type 2, 622
Type 1 diabetes, 45, 61, 182, 260, 279, 328, 596, 602, 673, 803, 815, 824, 887, 908, 1010, 1027, 1090, 1165, 1193, 1250, 1282
Type 2 diabetes, 61, 284, 293, 397, 460, 483, 526, 532, 649, 665, 708, 753, 766, 872, 900, 943, 974, 1003, 1033, 1048, 1175, 1212, 1242, 1255, 1282
Type 1 diabetes mellitus, 617, 961, 1135, 1235
Type 2 diabetes mellitus, 228, 466, 478, 510, 518, 589, 722, 791, 855, 864, 996, 1220, 1269, 1287
Type 1 glucose control, 740
Type 1b diabetes, 1070

UK, 803, 1269
ulcer care, 93
upper gastrointestinal haemorrhage, 70
urinary albumin excretion, 397
urinary pH, 556
utilization, 1105

variability, 714
vascular disease, 773
vascular endothelial growth factor, 247
vasomotion, 880
very low-density lipoprotein, 532
visual fields, 97
vitamin D_3, 19

waist circumference, 753
weight loss, 708
weight training, 220
white blood cell count, 955
wound healing, 89

X-PERT®, 40

yield, 1040
young adults, 354

© 2009 The Authors.
Journal compilation © 2009 Diabetes UK.